

Toric Varieties for Discrete Geometers

Francisco Santos, University of Cantabria

November 2020


Sources:

- B. Sturmfels, "Gröbner bases and convex polytopes",
Lect. Univ. Series, Amer. Math. Soc., 1995
- D. Cox, "What is a toric variety?"
in "Topics in algebraic geometry and geometric modeling"
Contemp. Math 334, Amer. Math. Soc., 2003, pp 203-223

(Plus: Fulton, "Introduction to toric varieties", Princeton U.P., 1993.

Cox, Little, Schenk, "Toric varieties", A.M.S., 2011.

The five ways to a toric variety



	affine	projective	abstract
normal	(*)	(IV)	(V)
perhaps not-normal	(I) (II)	(III)	

(*) (I) is normal $\Leftrightarrow A$ generates $\text{pos}(A) \cap \mathbb{Z}^d$

(II) is normal $\Leftrightarrow S = \sigma \cap \mathbb{Z}^d$ for a cone σ

(0) "Toric varieties for algebraic geometers"

Def: "A toric variety is a complex algebraic variety X that contains a torus $T \cong (\mathbb{C}^*)^d$ as an open dense subset, together with an action $T \cdot X \rightarrow X$ of T on X that extends the standard action (multiplication) $T \cdot T \rightarrow T$

Glossary:

Variety \rightarrow affine: the zero set of ^{a system of} finitely many polynomials $\subseteq \mathbb{C}^n$
 \rightarrow projective: same, but with homogeneous polynomials $\subseteq \mathbb{P}^n = \mathbb{P}(\mathbb{C}^{n+1})$
 \rightarrow abstract: (for this talk) an object constructed by gluing affine varieties via transition maps.

Normal: "soft version" of smooth. A local property, that limits the type of singularities that you allow.

Algebraic torus: $T^n = (\mathbb{C}^*)^n$, with coordinate-wise multiplication

Etymology: $(\mathbb{C}^*)^n \cong \underbrace{(\mathbb{S}^1)^n}_{\text{topologists' torus}} \times \underbrace{(0, \infty)^n}_{\text{topologically trivial}}$
as groups, with coordinate-wise multiplication, and $\mathbb{S}^1 = \{z \in \mathbb{C} \mid |z| = 1\}$


(0) "Toric varieties for combinatorial topologists":

"Cell complexes" made of tori, with a single top-dimensional torus plus lower tori "glued to it".

Example: \mathbb{C}^2

$$\mathbb{C}^2 = (\mathbb{C}^*)^2 \cup \begin{matrix} \mathbb{C}^* \times \{0\} \\ \{0\} \times \mathbb{C}^* \end{matrix} \cup \{0,0\}$$


\downarrow dim 2 $2 \times \downarrow$ dim 1 \downarrow dim 0



$\mathbb{C}^2 =$ "gluing of 4 tori, with f-vector (1, 2, 1)"

Example: \mathbb{P}^2

$$\mathbb{P}^2 = \{ (x, y, z) \in \mathbb{C}^3 \setminus \{0,0,0\} \} / \text{dilation}$$



$\mathbb{P}^2 =$ "gluing of 8 tori, with f-vector (3, 3, 1)"

(I) The toric variety of $A = \{a_1, \dots, a_m\} \subseteq \mathbb{Z}^d$

Embed $(\mathbb{C}^*)^d \hookrightarrow \mathbb{C}^m$ via A :

$$t = (t_1, \dots, t_d) \longmapsto (t^{a_1}, \dots, t^{a_m})$$

remark: negative exponents are ok since $t_i \neq 0$

Let $V_A := \text{closure}(\text{Im}((\mathbb{C}^*)^d))$

Remarks:

• linear relations in $A \longrightarrow$ binomials vanishing on V_A

• In fact: $I_A := I_{V(A)}$ is generated by those binomials


(sturmfels Prop. 4.11)

• $0 \in V_A \iff \text{pos}(A)$ is "pointed" or "strongly convex"
↳ special case of

• More generally:

THE "STRUCTURE THEOREM" OF TORIC VARIETIES

(my naming see appendix A.1)



(II) The toric variety of an affine semigroup

Semigroup of $A = \{a_1, \dots, a_m\}$: $S_A := \{n_1 a_1 + \dots + n_m a_m \mid n_i \in \mathbb{Z}_{\geq 0}\}$

Affine semigroup := S_A for some A

Remark: if $A \subseteq A' \subseteq S_A$, then
 $V_A \subseteq \mathbb{C}^m, V_{A'} \subseteq \mathbb{C}^{m'}, m' = |A'| > m$

new coords.
depend on old ones
polynomially

$V_A \cong V_{A'}$

forget coords

(adding positive combinations of existing points in A does not change the toric variety)

In particular, we can call V_A the "toric variety of S_A ".


Special (important) case: $S_\sigma = \sigma \cap \mathbb{Z}^d$ for a rat. polyhedral cone σ .

Gordon's Lemma: S_σ is finitely generated. $\rightarrow S_\sigma$ is an "affine semigroup"

It has a minimal generating set: the Hilbert basis of S_σ

In this case $V_\sigma := V_{\text{Hilbert}(\sigma)}$ is normal. In fact:

normal affine
toric varieties
(modulo algebraic
isomorphism)



[Thm 13.5, Sturmfels]
 rational polyhedral
cones in \mathbb{R}^d
 (modulo unimodular
integer transformation)

(III) The projective toric variety of a homogeneous A .

A is homogeneous := \exists lin. funct. f
 with $f(A) = \text{constant} \neq 0$
 Sturmfels, Ch 13,
 calls this "graded"

(that is: A is contained
 in an affine hyperplane
 away from 0)

In this case:

$\left. \begin{array}{l} \text{linear relations in } A \\ \text{(Sturmfels lemma 4.14)} \end{array} \right\} = \left. \begin{array}{l} \text{affine relations in } A \\ \text{affine geometry} \end{array} \right\}$

(linear algebra
 restricted to an
 affine hyperplane)

\Rightarrow the binomial ideal I_A is homogeneous

\Rightarrow the affine variety $V_A \subseteq \mathbb{C}^m$ is a union of lines

\Rightarrow it gives a projective variety \tilde{V}_A in \mathbb{P}^{m-1}

For homogeneous A :

$\left. \begin{array}{l} \text{poset of tori} \\ \text{in } \tilde{V}_A \end{array} \right\} = \left. \begin{array}{l} \text{poset of} \\ \text{faces of} \\ \text{conv}(A) \end{array} \right\}$

poset of "tori" of V_A ,
 excluding 0 and
 reducing dimensions by 1

poset of "faces" of $\text{pos}(A)$,
 excluding 0 and
 reducing dimensions by 1

(IV) The projective toric variety of a polytope.

Now let P be a ^{lattice} polytope in \mathbb{R}^d .

We can build the projective toric variety of $P \cap \mathbb{Z}^d$, but this is not what people call "toric variety of P ".

Remark: if we dilate P by an integer factor $\nu \in \mathbb{N}$, at some point $\widetilde{V}_{\nu P \cap \mathbb{Z}^d}$ becomes normal and after that it does not change (modulo isomorphism).

We define

$$X_P = \widetilde{V}_{\nu P \cap \mathbb{Z}^d}, \text{ for } \nu \text{ "large enough"}$$

($\nu = d$ is enough)

Remark: affine charts
in X_P



vertex (or face)
figures in P

Remark: If P and P' are normally equivalent \rightarrow same normal fan then $X_P \cong X_{P'}$ because the construction depends only on the "vertex cones" $\{ \text{pos}(P - v) \mid v \text{ a vertex of } P \}$

(V) The toric variety of a fan

Let \mathcal{N} be a rat. polyh. fan in \mathbb{R}^d (E.g. the normal fan of a P)

① For each cone $\sigma \in \mathcal{N}$ (i.e. face F of P) consider

the polar cone σ^\vee (that is, the face cone $\text{pos}(P-F)$)

and build the affine normal toric variety $U_\sigma := V_{\sigma^\vee} \cap \mathbb{Z}^d$

(the affine chart of the X_P in the previous slide).

② Glue these affine charts in the natural way
(e.g. they all contain one and the same torus $T \cong (\mathbb{C}^*)^d$)

Remark:

- this recipe works for arbitrary fans.
- all normal toric varieties (in the sense of def 3) can be constructed in this way.

<u>Fan</u>	<u>Variety</u>
complete	complete (\Leftrightarrow compact)
polytopal	projective
"regular"	quasi-projective.

(A.1) The structure theorem

Let $A = \{a_1, \dots, a_m\} \subseteq \mathbb{Z}^d$. Let $C \subseteq [m]$.

Notation:

1) $x|_C =$ "x restricted to C", for $x \in \mathbb{C}^m$

2) $1_C =$ "the 0/1 vector with support C"

T. F. A. E:

① $\exists c: \mathbb{R}^d \rightarrow \mathbb{R}$ with $c(a_i) \begin{cases} = 0 & \text{if } i \notin C \\ > 0 & \text{if } i \in C \end{cases}$ "positive covectors"
"faces of pos(A)"

② \nexists linear dependence $\sum_{i=1}^m \lambda_i a_i = 0$ with $\lambda_i \geq 0 \forall i$ and $\lambda_i = 0$ if $i \notin C$ positive dependencies

③ $1_{[n] \setminus C} \in V_A$

④ V_A contains an x with $\text{supp}(x) = [n] \setminus C$

proof: ① \Leftrightarrow ② is Farkas Lemma (or, "vector/covector orthogonality")

③ \Rightarrow ④ is trivial

① \Rightarrow ③ Let c be as in ① and

notice that:

$$\lim_{t \rightarrow 0} (t^{c a_1}, \dots, t^{c a_m}) = 1_{[n] \setminus C}$$

As t varies in \mathbb{C}^* this is a "1-parameter subgroup" in $\text{Im}(\phi)$.

④ \Rightarrow ② Suppose ② fails, and let

λ be as in ②. Since $\lambda \geq 0$, its binomial is of the form $x^\lambda - 1 = 0$

In particular, $\forall x = (x_1, \dots, x_m) \in V_A$

$$|x_1|^{\lambda_1} \cdot \dots \cdot |x_m|^{\lambda_m} = 1$$

\Rightarrow some x_i with $i \in C$ has $|x_i| \geq 1$

\Rightarrow No $x \in V_A$ has support $\subseteq [n] \setminus C$ \square

That is: $\text{faces of pos(A)} = \text{stratification of } V_A \text{ by support} = \text{0/1 points in } V_A$
 (Cox, Sect. 9) as posets, and as labelled by their C \leftarrow strata are the orbits of the $(\mathbb{C}^*)^d$ action, and they are themselves tori \leftarrow limit points of 1-parameter subgroups

for "abstract" $t.v.$, via fans

(A.2) The local structure (affine charts) of X_P

For any homogeneous A we have

$$\begin{array}{ccc} (\mathbb{C}^*)^d & \xrightarrow{\phi} & \text{Im}(\phi) \subseteq V_A & \longrightarrow & V_A \\ t & \longmapsto & (t^{a_1}, \dots, t^{a_m}) & \text{quotient } \mathbb{P}^{m-1} = \mathbb{C}^m / \sim \end{array}$$

Each of the m coordinates in \mathbb{P}^{m-1} gives us an affine chart, obtained by setting it = 1.

That is, the i -th chart is the affine t.o.v.

$$\text{of } A\text{-}\{a_i\} : \left(\begin{array}{ccc} (\mathbb{C}^*)^d & \xrightarrow{\phi_i} & \text{Im}(\phi_i) \subseteq V_{A\text{-}\{a_i\}} = \overline{\text{Im}(\phi_i)} \\ t & \longmapsto & (t^{a_1 - a_i}, \dots, 1, \dots, t^{a_m - a_i}) \end{array} \right)$$

That is :

(Sturmfels Lemma 13.10)

Affine charts of

$$V_A$$

polytope language

Affine toric varieties
of the "vertex figures" (*)
or "contractions" of A

oriented matroid language

(*) strictly speaking, we have one chart for each lattice point in P , not only vertices

In particular:

As we dilate P , the affine charts depend on more and more points in the vertex cones.

After we get a Hilbert basis in every cone, the affine charts change no more.



On the other hand, since normality is local, when all affine charts are normal so is V_A

In particular, to say that $V_{\mathbb{P}^n/\mathbb{Z}^d}$ stabilizes we need to look at how the charts of non-vertices glue to those of vertices, but...