

Constructible sheaves and functions up to infinity

Pierre Schapira

April 22, 2021

Abstract

We introduce the category of b-analytic manifolds, a natural tool to define constructible sheaves and functions up to infinity. We study with some details the operations on these objects and also recall the Radon transform for constructible functions.

Contents

1	A short review on sheaves	4
2	Constructible sheaves up to infinity	7
2.1	Subanalytic subsets up to infinity	7
2.2	Constructible sheaves up to infinity	9
2.3	Convolution and γ -topology	11
3	A short review on constructible functions	13
3.1	From constructible sheaves to constructible functions	13
3.2	Operations	15
4	Constructible functions up to infinity	17
4.1	Definition	17
4.2	Operations	19
4.3	γ -constructible functions	22
5	Correspondences for constructible functions	23
5.1	Correspondences	23
5.2	Application: the Radon transform	25

Key words: constructible sheaves, constructible functions, subanalytic geometry

MSC: 55N99, 32B20, 32S60

This research was supported by the ANR-15-CE40-0007 “MICROLOCAL”.

Introduction

The triangulated category of constructible sheaves over a commutative Noetherian ring \mathbf{k} on a real analytic manifold plays an increasing role in various fields of mathematics and is perfectly understood, as well as its Grothendieck group which is known to be isomorphic (when \mathbf{k} is a field of characteristic 0) to the group of constructible functions as well as to that of Lagrangian cycles.

However, it often happens that one is led to consider such objects “up to infinity”. Although this point is certainly not totally new (see below), no systematic study exists to our knowledge and the aim of this paper is to fill this gap. As we shall see, to consider constructibility up to infinity is much more natural and makes several hypotheses of properness useless.

Recall that constructible functions and Lagrangian cycles first appeared in the algebraic setting with Robert McPherson [McP74] and in the complex analytic setting with Masaki Kashiwara [Kas73]. In the complex setting, Lagrangian cycles were studied by several people and in particular by Victor Ginsburg [Gin86] and Claude Sabbah [Sab85] for their functorial properties. The real case was first treated in [Kas85]. See also [KS90, Ch. IX, Notes] for an history of the subject.

The Euler calculus of constructible functions has been introduced independently by Oleg Viro (see [Vir88]) in the complex analytic setting and by the author in the subanalytic setting (see [Sch89]). It has many applications, particularly to tomography *i.e.*, real Radon transform, see [Sch95] (see also Lars Ernström [Ern94] for complex projective duality) and more generally in Topological Data Analysis (see [CGR12] for a survey).

Lagrangian cycles are not so easy to describe, contrarily to constructible functions and we shall not study them here.

A constructible function φ on a real analytic manifold X is mathematically very simple: it is a \mathbb{Z} -valued function, the sets $\varphi^{-1}(m)$ ($m \in \mathbb{Z}$) being all subanalytic and the family of such sets being locally finite. It is not difficult (with the tools of subanalytic geometry at hands) to check that the set $\mathcal{CF}(X)$ of constructible functions on X is a commutative unital algebra and that the inverse image (*i.e.*, composition) of such a function by a morphism $f: Z \rightarrow X$ is again constructible. Things become more unusual when looking at direct images, in particular integration. Assume that φ has compact support. One may write φ as a finite sum $\sum_{i \in I} c_i \mathbf{1}_{K_i}$ where $c_i \in \mathbb{Z}$, K_i is a compact subanalytic subset of X and for $S \subset X$, $\mathbf{1}_S$ is the characteristic function of S . Then one defines the integral of φ by the formula

$$\int_X \varphi = \sum_{i \in I} c_i \cdot \chi(K_i)$$

where $\chi(K_i)$ denotes the Euler-Poincaré index of K_i . (Of course, one has to check that this definition does not depend on the decomposition of φ .) For a morphism $f: X \rightarrow Y$ of real analytic manifolds, one defines the integral along f of a function $\varphi \in \mathcal{CF}(X)$ whose support is proper with respect to f by setting for $y \in Y$,

$$\left(\int_f \varphi\right)(y) = \int_X \varphi \cdot \mathbf{1}_{f^{-1}(y)},$$

and one checks that one obtains a constructible function on Y . This integral has all properties of classical integrals (linearity and Fubini theorem, that is, functoriality), except that it is not positive (the integral of $\mathbf{1}_{(0,1)}$ is -1) and a set reduced to one point has integral 1. In fact, one easily translates all operations on constructible sheaves to operations on constructible functions. In particular duality makes sense for constructible functions and commutes with direct images.

As mentioned in the title, we shall extend all these constructions to the case of constructible sheaves and functions “up to infinity”. For that purpose we introduce the category of b-analytic manifolds. An object X_∞ is an open embedding $X \subset \widehat{X}$ of smooth real analytic manifolds with X subanalytic and relatively compact in \widehat{X} , and a morphism $f: X_\infty \rightarrow Y_\infty$ is a real analytic map $f: X \rightarrow Y$ such that the graph of f is subanalytic in $\widehat{X} \times \widehat{Y}$. This definition has a certain similarity with that of bordered space of [DK16] but is different.

Note that the notion of being subanalytic up to infinity is closely related to that of definable sets and of \mathcal{o} -minimal structures, well known from the specialists (see in particular [VdD98, VdDM96]) and constructible sheaves and functions in this framework have already been defined in [Sch03, EP20]. Nevertheless, our approach for sheaves, based on the notion of micro-support, is of a different nature.

Sections 1 and 3 are short reviews on sheaves and constructible functions, posted here for the reader’s convenience.

In **Section 2** we define a (derived) sheaf constructible up to infinity on X as a constructible sheaf whose micro-support is subanalytic in the cotangent bundle $T^*\widehat{X}$. This is equivalent to saying that its (proper or non proper) direct image in \widehat{X} is again constructible. Note that such a property already appeared in [KS20]. We briefly study the six operations on the triangulated category of derived constructible sheaves up to infinity. Contrarily to the classical constructible case, the two inverse images f^{-1} and $f^!$ are exchanged by duality, the two direct images Rf_* and $Rf_!$ are constructible without any properness assumptions and, again, are exchanged by duality. As a nice application, we find that non proper convolution on a real vector space \mathbb{V} is well defined on constructible sheaves up to infinity and is associative. Such a non proper convolution appears when using the so-called γ -topology, associated with a closed convex proper cone γ of \mathbb{V} .

In **Section 4**, we define the space $\mathcal{CF}(X_\infty)$ of constructible functions up to infinity and study with some care the operations on such functions. Contrarily to the classical case, we have now two kind of integrals, proper and non proper, and, as for sheaves, these operations are defined without any properness hypothesis. Moreover, they are exchanged by duality.

In **Section 5**, posted here for easier accessibility, we recall (and adapt) the main results of [Sch95] in which we obtain an inversion formula for the Radon transform of constructible functions. This formula asserts that one can recover a constructible function on a real vector space \mathbb{V} from the knowledge of the Euler-Poincaré index of its restriction to all affine hyperplanes. For example, if $\dim \mathbb{V} = 3$, one can reconstruct a compact subanalytic subset from the knowledge of the number of connected components and holes of the restriction of the compact set to all slices (affine planes).

To summarize, it appears, and it is not surprising, that the notion of being constructible up to infinity is much more natural than the usual notion.

Acknowledgement. I warmly thank Ezra Miller for several fruitful comments on a previous version of this paper as well as François Petit for several stimulating discussions.

Convention. In this paper, \mathbf{k} denotes a commutative unital Noetherian ring. From Section 3 and until the end of the paper, \mathbf{k} is a field of characteristic 0.

1 A short review on sheaves

Recall that a topological space is *good* if it is Hausdorff, locally compact, countable at infinity and of finite flabby dimension.

For a space X , we denote by Δ_X the diagonal of $X \times X$ and if $f: X \rightarrow Y$ is a map, we denote by Γ_f its graph in $X \times Y$. We denote by pt the space consisting of a single point and by $a_X: X \rightarrow \text{pt}$ the unique map from X to pt .

We consider a commutative unital Noetherian ring \mathbf{k} and a good topological space X . We denote by $D^b(\mathbf{k}_X)$ the bounded derived category of sheaves of \mathbf{k} -modules on X and simply call an object of this category “a sheaf”. We shall freely make use of the six Grothendieck operations on sheaves and refer to [KS90] for an exposition and for notations. In particular, we denote by ω_X the dualizing complex, $\omega_X := a_X^! \mathbf{k}_{\{\text{pt}\}}$. We have the duality functors

$$D'_X(\bullet) = R\mathcal{H}om(\bullet, \mathbf{k}_X), \quad D_X = R\mathcal{H}om(\bullet, \omega_X).$$

Consider a locally closed subset Z of X and denote by $j_Z: Z \hookrightarrow X$ the embedding. One sets $\mathbf{k}_{XZ} := j_{Z!} \mathbf{k}_Z$. This is the sheaf on X which is the constant sheaf with stalk \mathbf{k} on Z and 0 elsewhere. If Z is closed, $j_{Z!} \mathbf{k}_Z \simeq j_{Z*} \mathbf{k}_Z$. If there is no risk of confusion (in particular when Z is closed), we write \mathbf{k}_Z instead of \mathbf{k}_{XZ} . If F is a sheaf on X , we set $F_Z := F \otimes \mathbf{k}_{XZ}$.

Kernels

Given topological spaces X_i ($i = 1, 2, 3$) we set $X_{ij} = X_i \times X_j$, $X_{123} = X_1 \times X_2 \times X_3$. We denote by $q_i: X_{ij} \rightarrow X_i$ and $q_{ij}: X_{123} \rightarrow X_{ij}$ the projections.

$$(1.1) \quad \begin{array}{ccc} & X_{12} & \\ q_1 \swarrow & & \searrow q_2 \\ X_1 & & X_2 \end{array} \quad \begin{array}{ccc} & X_{123} & \\ q_{12} \swarrow & \downarrow q_{13} & \searrow q_{23} \\ X_{12} & X_{13} & X_{23} \end{array}$$

For $A \subset X_{12}$ and $B \subset X_{23}$ one sets

$$(1.2) \quad A \times_2 B = A \times_{X_2} B = q_{12}^{-1} A \cap q_{23}^{-1} B, \quad A \circ_2 B = q_{13}(A \times_2 B).$$

For good topological spaces X_i 's as above, one often calls an object $K_{ij} \in D^b(\mathbf{k}_{X_{ij}})$ a *kernel*. One defines as usual the composition of kernels

$$(1.3) \quad K_{12} \circ_2 K_{23} := Rq_{13!}(q_{12}^{-1}K_{12} \overset{L}{\otimes} q_{23}^{-1}K_{23}).$$

If there is no risk of confusion, we write \circ instead of \circ_2 .

It is easily checked, and well known, that convolution is associative, namely given three kernels $K_{ij} \in D^b(\mathbf{k}_{X_{ij}})$, $i = 1, 2, 3$, $j = i + 1$ one has an isomorphism

$$(1.4) \quad (K_{12} \circ K_{23}) \circ K_{34} \simeq K_{12} \circ (K_{23} \circ K_{34}),$$

this isomorphism satisfying natural compatibility conditions that we shall not make here explicit.

Of course, this construction applies in the particular case where $X_i = \text{pt}$ for some i . For example, if $K \in D^b(\mathbf{k}_{X \times Y})$ and $F \in D^b(\mathbf{k}_X)$, one usually sets $\Phi_K(F) = F \circ K$. Hence

$$(1.5) \quad \Phi_K(F) = F \circ K = Rq_{2!}(q_1^{-1}F \overset{L}{\otimes} K).$$

We shall also use the right adjoint of the functor $\Phi_K(\cdot)$, namely the functor $\Psi_K(\cdot)$ (see [KS90, § 3.6]), defined for $G \in D^b(\mathbf{k}_Y)$ by:

$$(1.6) \quad \Psi_K(G) = Rq_{1*}R\mathcal{H}om(K, q_2^!G).$$

The next result is elementary. Denote by K^v the image of $K \in D^b(\mathbf{k}_{X \times Y})$ by the map $X \times Y \xrightarrow{\sim} Y \times X$, $(x, y) \mapsto (y, x)$.

Lemma 1.1. *Let $f: X \rightarrow Y$, $F \in D^b(\mathbf{k}_X)$ and $G \in D^b(\mathbf{k}_Y)$. Set for short $K_f = \mathbf{k}_{\Gamma_f}$. Then*

$$\begin{aligned} f^{-1}G &\simeq K_f \circ G = \Phi_{K_f^v}G, & Rf_*F &\simeq Rq_{2*}R\mathcal{H}om(K_f, q_1^!F) = \Psi_{K_f^v}F, \\ Rf_!F &\simeq F \circ K_f = \Phi_{K_f}F, & f^!G &\simeq Rq_{1*}R\mathcal{H}om(K_f, q_2^!G) = \Psi_{K_f}(G). \end{aligned}$$

Proof. The first and third isomorphisms are obvious (identify X with Γ_f). The two others follow by adjunction. \square

Remark 1.2. Let $K \in D^b(\mathbf{k}_{X \times Y})$, $F \in D^b(\mathbf{k}_X)$ and $G \in D^b(\mathbf{k}_Y)$. Set

$$F \overset{\text{np}}{\circ} K = Rq_{2*}(q_1^{-1}F \overset{L}{\otimes} K), \quad K \overset{\text{np}}{\circ} G = Rq_{1*}(K \overset{L}{\otimes} q_2^{-1}G).$$

Denote by $j: \Gamma_f \hookrightarrow X \times Y$ the embedding of the graph of f . One has

$$\begin{aligned} Rf_*F &\simeq Rq_{2*}R\mathcal{H}om(K_f, q_1^!F) \simeq Rq_{2*}j_!j^!q_1^!F \\ &\simeq Rq_{2*}j_!j^{-1}q_1^{-1}F \simeq Rq_{2*}(q_1^{-1}F \overset{L}{\otimes} \mathbf{k}_{\Gamma_f}) \simeq F \overset{\text{np}}{\circ} K_f. \end{aligned}$$

Micro-support

Now assume that X is a real manifold of class C^∞ and denote by $\pi_X: T^*X \rightarrow X$ its cotangent bundle. To $F \in D^b(\mathbf{k}_X)$, one associates its *micro-support* $\text{SS}(F)$ (also called *singular support*), a closed \mathbb{R}^+ -conic subset of T^*X and this set is co-isotropic (in a sense that we do not recall here). See [KS90, Th. 6.5.4].

Subanalytic subsets

From now on and unless otherwise specified, we work on real analytic manifolds. However, almost all results extend to the case of subanalytic spaces for the definition of which we refer to [KS16, § 2.4].

We shall not review here the history of subanalytic geometry, which takes its origin in the work of Lojasiewicz, simply mentioning the names of Gabrielov and Hironaka. References are made to [BM88].

Let X be a real analytic manifold. Denote by \mathcal{S}_X the family of subanalytic subsets of X . Then \mathcal{S}_X is a Boolean algebra which contains the family of semi-analytic subsets (those locally defined by analytic inequalities) and is stable by taking the closure and the interior. If $f: X \rightarrow Y$ is subanalytic, $A \in \mathcal{S}_X$, $B \in \mathcal{S}_Y$, then $f^{-1}(B) \in \mathcal{S}_X$ and if f is proper on the closure of A , then $f(A) \in \mathcal{S}_Y$.

Moreover, to be subanalytic in X is a local property on X . More precisely, given $X = \bigcup_{a \in A} U_a$ an open covering, a subset $Z \subset X$ is subanalytic in X if and only if $Z \cap U_a$ is subanalytic in U_a for all $a \in A$.

Note that if Z is a locally closed subanalytic subset of X , then there exist an open set U and a closed subset S both subanalytic in X such that $Z = U \cap S$. Indeed, set $Y = \overline{Z} \setminus Z$. Then Y is closed since Z is locally closed. Choose $S = \overline{Z}$ and $U = X \setminus Y$.

A subanalytic stratification of X is a locally finite stratification $X = \bigsqcup_{a \in A} X_a$ where each X_a is a smooth locally closed real analytic submanifold of X , subanalytic in X .

Constructible sheaves

A sheaf $F \in D^b(\mathbf{k}_X)$ is weakly \mathbb{R} -constructible if there exists a subanalytic stratification $X = \bigsqcup_{a \in A} X_a$ such that for all $j \in \mathbb{Z}$, $H^j(F)|_{X_a}$ is locally constant. If moreover, these locally constant sheaves are of finite rank, then F is \mathbb{R} -constructible. By the results of [KS90, Ch. VIII], F is weakly \mathbb{R} -constructible if and only if $\text{SS}(F)$ is contained in a closed conic subanalytic isotropic subvariety of T^*X and this implies that $\text{SS}(F)$ is equal to a closed conic subanalytic Lagrangian subvariety.

One denotes by $D_{\mathbb{R}c}^b(\mathbf{k}_X)$ the full triangulated subcategory of $D^b(\mathbf{k}_X)$ consisting of \mathbb{R} -constructible sheaves. The categories of constructible sheaves are stable by the six Grothendieck operations with the exception of direct images which should be proper on the supports of the constructible sheaves.

2 Constructible sheaves up to infinity

2.1 Subanalytic subsets up to infinity

In order to define subanalytic subsets up to infinity, we introduce the category of b-analytic manifolds, inspired by (but rather different from) that of bordered space of [DK16]. As mentioned in the introduction, the notion of being subanalytic up to infinity is a particular case of that of definable set, well known from the specialists (see [VdD98, VdDM96]), and constructible sheaves in this framework have already been defined in [Sch03, EP20]. However, our approach is direct and quite different since it is based on the notion of micro-support.

Definition 2.1. The category of *b-analytic manifolds* is the category defined as follows.

- (a) An object X_∞ is a pair (X, \widehat{X}) with $X \subset \widehat{X}$ an open embedding of real analytic manifolds such that X is relatively compact and subanalytic in \widehat{X} ,
- (b) a morphism $f: X_\infty = (X, \widehat{X}) \rightarrow Y_\infty = (Y, \widehat{Y})$ of b-analytic manifolds is a morphism of real analytic manifolds $f: X \rightarrow Y$ such that the graph Γ_f of f in $X \times Y$ is subanalytic in $\widehat{X} \times \widehat{Y}$.
- (c) The composition $(X, \widehat{X}) \xrightarrow{f} (Y, \widehat{Y}) \xrightarrow{g} (Z, \widehat{Z})$ is given by $g \circ f: X \rightarrow Z$ and the identity $\text{id}_{(X, \widehat{X})}$ is given by id_X (see Lemma 2.3 below).

If there is no risk of confusion, we shall often denote by $j_X: X \hookrightarrow \widehat{X}$ the open embedding.

Remark 2.2. Instead of requiring \widehat{X} to be a smooth real analytic manifold and X relatively compact in it, one could ask \widehat{X} to be a compact subanalytic space in the sense of [KS16, § 2.4]. However, Definition 2.8 should be modified. One could also define the notion of a b-subanalytic space.

Remark that in general, contrarily to the case of bordered spaces, neither (X, X) nor $(\widehat{X}, \widehat{X})$ are b-analytic manifolds. However, if X is compact, (X, X) is a b-analytic manifold.

Lemma 2.3. (a) *The identity $\text{id}_{(X, \widehat{X})}$ is a morphism of b-analytic manifolds.*

(b) *Let $f: (X, \widehat{X}) \rightarrow (Y, \widehat{Y})$ and $g: (Y, \widehat{Y}) \rightarrow (Z, \widehat{Z})$ be morphisms of b-analytic manifolds. Then the composition $g \circ f$ is a morphism of b-analytic manifolds.*

Proof. (a) Since X is subanalytic in \widehat{X} , $X \times X$ is subanalytic in $\widehat{X} \times \widehat{X}$, and $\Delta_X = X \times X \cap \Delta_{\widehat{X}}$ is subanalytic in $\widehat{X} \times \widehat{X}$.

(b) By the hypothesis, Γ_g is subanalytic and relatively compact in $\widehat{Y} \times \widehat{Z}$ and Γ_f is subanalytic and relatively compact in $\widehat{X} \times \widehat{Y}$. It follows that $\Gamma_f \times_{\widehat{Y}} \Gamma_g$ is subanalytic and relatively compact in $\widehat{X} \times \widehat{Y} \times \widehat{Z}$. Therefore, its projection $\Gamma_f \circ \Gamma_g$ is subanalytic in $\widehat{X} \times \widehat{Z}$. Since $\Gamma_f \circ \Gamma_g = \Gamma_{g \circ f}$, the proof is complete. (Note that one could also have applied Proposition 2.6 below.) \square

Definition 2.4. Let $X_\infty = (X, \widehat{X})$ be a b-analytic manifold and let Z be a subset of X . We say that Z is subanalytic in X_∞ if Z is subanalytic in \widehat{X} . If there is no risk of confusion (that is, \widehat{X} has been already defined) we also say that Z is subanalytic up to infinity.

Note that the family of subsets subanalytic up to infinity inherits of all properties of the family of subanalytic subsets with the exception that this property is no more local. In particular, this family is stable by closure, interior, complement, finite unions and finite intersections and X itself is subanalytic up to infinity.

On a real analytic manifold X , the subanalytic topology and the site X_{sa} are defined in [KS01].

Definition 2.5. Let $X_\infty = (X, \widehat{X})$ be a b-analytic manifold.

- (a) We shall denote by $\text{Op}_{X_{\infty\text{sa}}}$ the category of open subsets of X subanalytic up to infinity, the morphisms being the inclusions.
- (b) We endow $\text{Op}_{X_{\infty\text{sa}}}$ with a Grothendieck topology as follows. A family $\{U_i\}_{i \in I}$ of objects of $\text{Op}_{X_{\infty\text{sa}}}$ is a covering of $U \in \text{Op}_{X_{\infty\text{sa}}}$ if $U_i \subset U$ for all $i \in I$ and there exists $J \subset I$ with J finite such that $U = \bigcup_{j \in J} U_j$.
- (c) We denote by $X_{\infty\text{sa}}$ the site so obtained.

Note that the category $\text{Op}_{X_{\infty\text{sa}}}$ is stable by product of two elements (namely, the intersection of two open subsets) and admits a terminal object, namely X . This makes the study of sheaves on $X_{\infty\text{sa}}$ particularly easy.

In the sequel, for $U \in \text{Op}_{X_{\infty\text{sa}}}$, we shall denote by U_∞ the b-analytic manifold (U, \widehat{X}) where the embedding $j_U: U \hookrightarrow \widehat{X}$ is the composition of j_X and the embedding $U \hookrightarrow X$.

Proposition 2.6. Let $X_{i\infty} = (X_i, \widehat{X}_i)$ ($i = 1, 2, 3$) be three b-analytic manifolds.

- (a) Setting $\widehat{X}_{12} = \widehat{X}_1 \times \widehat{X}_2$, the pair $(X_{12}, \widehat{X}_{12})$ is a b-analytic manifold. Moreover, if S_1 and S_2 are two subsets of X_1 and X_2 respectively, subanalytic up to infinity, then $S_1 \times S_2$ is subanalytic up to infinity in X_{12} .
- (b) Let S_1 and S_2 be subanalytic up to infinity in X_{12} and X_{23} respectively, then $S_1 \circ_2 S_2$ is subanalytic up to infinity in X_{13} .
- (c) In particular, let $f: X_\infty \rightarrow Y_\infty$ be a morphism of b-analytic manifolds. If $Z \subset Y$ is subanalytic up to infinity, then $f^{-1}(Z)$ is subanalytic up to infinity in X and if $S \subset X$ is subanalytic up to infinity, then $f(S)$ is subanalytic up to infinity in Y .

We shall denote by $(X \times Y)_\infty$ the b-analytic manifold $(X \times Y, \widehat{X} \times \widehat{Y})$.

Proof. (a) is obvious.

(b) $S_1 \times_{X_2} S_2$ is subanalytic and relatively compact in \widehat{X}_{123} . Therefore, its image by q_{13} is subanalytic and relatively compact in \widehat{X}_{13} .

(c) By the hypothesis, Γ_f is subanalytic up to infinity in $\widehat{X} \times \widehat{Y}$. By (b), $f^{-1}(Z) = \Gamma_f \circ_Y Z$ is subanalytic up to infinity in X and $f(S) = S \circ_X \Gamma_f$ is subanalytic up to infinity in Y . \square

2.2 Constructible sheaves up to infinity

In this section, we consider b-analytic manifolds $X_\infty = (X, \widehat{X})$ and $Y_\infty = (Y, \widehat{Y})$.

Definition

Let $F \in D_{\mathbb{R}c}^b(\mathbf{k}_X)$. Recall that the micro-support $\text{SS}(F)$ of F is a closed \mathbb{R}^+ -conic subanalytic Lagrangian subset of T^*X .

Lemma 2.7 (See [KS20, Th.2.2]). *Let $F \in D_{\mathbb{R}c}^b(\mathbf{k}_X)$. The following conditions are equivalent.*

- (a) *The micro-support $\text{SS}(F)$ is subanalytic in $T^*\widehat{X}$.*
- (b) *The micro-support $\text{SS}(F)$ is contained in a locally closed \mathbb{R}^+ -conic subanalytic isotropic subset of $T^*\widehat{X}$.*
- (c) *$j_{X!}F \in D_{\mathbb{R}c}^b(\mathbf{k}_{\widehat{X}})$.*
- (d) *$Rj_{X*}F \in D_{\mathbb{R}c}^b(\mathbf{k}_{\widehat{X}})$.*

Proof. For the reader's convenience, we recall the proof of loc. cit. Note that in loc. cit. the statement was formulated slightly differently.

(a) \Rightarrow (b) is obvious.

(c) \Rightarrow (a) and (d) \Rightarrow (a) follow from the fact that T^*X is subanalytic in $T^*\widehat{X}$. Indeed, set either $\Lambda = \text{SS}(j_{X!}F)$ or $\Lambda = \text{SS}(Rj_{X*}F)$. Then Λ is subanalytic in $T^*\widehat{X}$ and $\text{SS}(F) = \Lambda \cap T^*X$ is still subanalytic in $T^*\widehat{X}$.

(b) \Rightarrow (c). Assume that $\text{SS}(F)$ is contained in a locally closed \mathbb{R}^+ -conic subanalytic isotropic subset Λ of $T^*\widehat{X}$. By [KS90, Cor. 8.3.22], there exists a μ -stratification $\widehat{X} = \bigsqcup_{a \in A} Y_a$ such that $\Lambda \subset \bigsqcup_{a \in A} T_{Y_a}^*\widehat{X}$.

Set $X_a = X \cap Y_a$. Then $X = \bigsqcup_{a \in A} X_a$ is a μ -stratification and one can apply loc. cit. Prop. 8.4.1. Hence, for each $a \in A$, $F|_{X_a}$ is locally constant of finite rank. Hence $(j_{X!}F)|_{X_a}$ as well as $(j_{X!}F)_{\widehat{X} \setminus X} \simeq 0$ is locally constant of finite rank. Hence $j_{X!}F \in D_{\mathbb{R}c}^b(\mathbf{k}_{\widehat{X}})$.

(iv) (b) \Rightarrow (d). Set $G = j_{X!}F$. Then $G \in D_{\mathbb{R}c}^b(\mathbf{k}_{\widehat{X}})$ by (iii) and so does $Rj_{X*}F \simeq R\mathcal{H}om(\mathbf{k}_X, G)$ (apply [KS90, Prop. 8.4.10]). \square

Definition 2.8. Let $F \in D_{\mathbb{R}c}^b(\mathbf{k}_X)$. One says that F is constructible up to infinity if it satisfies one of the equivalent conditions in Lemma 2.7. We denote by $D_{\mathbb{R}c}^b(\mathbf{k}_{X_\infty})$ the full triangulated subcategory of $D_{\mathbb{R}c}^b(\mathbf{k}_X)$ consisting of sheaves constructible up to infinity.

It follows that if $F \in D_{\mathbb{R}c}^b(\mathbf{k}_{\widehat{X}})$, then $j_X^{-1}F \in D_{\mathbb{R}c}^b(\mathbf{k}_{X_\infty})$.

Example 2.9. Piecewise linear sheaves (PL sheaves) on a real vector space \mathbb{V} are defined in [KS19, Def. 2.3]. Clearly, PL-sheaves are constructible up to infinity.

Operations

Proposition 2.10. *Let X_∞ and Y_∞ be two b -analytic manifolds.*

- (i) *Let $F \in D_{\mathbb{R}c}^b(\mathbf{k}_{X_\infty})$ and $G \in D_{\mathbb{R}c}^b(\mathbf{k}_{Y_\infty})$. Then $F \boxtimes^L G \in D_{\mathbb{R}c}^b(\mathbf{k}_{(X \times Y)_\infty})$.*
- (ii) *Let F_1 and F_2 belong to $D_{\mathbb{R}c}^b(\mathbf{k}_{X_\infty})$. Then $F_1 \otimes^L F_2$ and $R\mathcal{H}om(F_1, F_2)$ belong to $D_{\mathbb{R}c}^b(\mathbf{k}_{X_\infty})$. In particular, the dual $D_X F$ of $F \in D_{\mathbb{R}c}^b(\mathbf{k}_{X_\infty})$ belongs to $D_{\mathbb{R}c}^b(\mathbf{k}_{X_\infty})$.*

Proof. (i) One has $\text{SS}(F \boxtimes^L G) \subset \Lambda_1 \times \Lambda_2$ with Λ_i an \mathbb{R}^+ -conic subset of T^*X_i subanalytic in $T^*\widehat{X}_i$ and isotropic ($i = 1, 2$). Then $\Lambda_1 \times \Lambda_2$ has the same property in $T^*(\widehat{X}_1 \times \widehat{X}_2)$. (One could also use the fact that $j_{X \times Y!}(F \boxtimes G) \simeq j_{X!}F \boxtimes j_{Y!}G$.)

(ii) Denote by Λ_i the micro-support of F_i , $i = 1, 2$. By [KS90, Cor. 6.4.5], the micro-support of $F_1 \otimes^L F_2$ is contained in $\Lambda_1 \hat{+} \Lambda_2$ and this set is conic subanalytic isotropic in \widehat{X} by [KS90, Cor. 8.3.18 (i)]. The proof for $R\mathcal{H}om$ is similar. \square

Proposition 2.11. *Let $f: X_\infty \rightarrow Y_\infty$ be a morphism of b -analytic manifolds.*

- (i) *Let $G \in D_{\mathbb{R}c}^b(\mathbf{k}_{Y_\infty})$. Then $f^{-1}(G)$ and $f^!G$ belong to $D_{\mathbb{R}c}^b(\mathbf{k}_{X_\infty})$.*
- (ii) *Let $F \in D_{\mathbb{R}c}^b(\mathbf{k}_{X_\infty})$. Then $Rf_!F$ and Rf_*F belong to $D_{\mathbb{R}c}^b(\mathbf{k}_{Y_\infty})$.*

Proof. Let $K_f = \mathbf{k}_{\Gamma_f}$. Then $K_f \in D_{\mathbb{R}c}^b(\mathbf{k}_{(X \times Y)_\infty})$. By Proposition 2.10 and Lemma 1.1, we are reduced to prove that

- (a) if $H \in D_{\mathbb{R}c}^b(\mathbf{k}_{(X \times Y)_\infty})$, then $Rq_{1!}H$ and $Rq_{1*}H$ belong to $D_{\mathbb{R}c}^b(\mathbf{k}_{X_\infty})$,
- (b) if $F \in D_{\mathbb{R}c}^b(\mathbf{k}_{X_\infty})$, then $q_1^{-1}F$ and $q_1^!F$ belong to $D_{\mathbb{R}c}^b(\mathbf{k}_{(X \times Y)_\infty})$.

The assertion (b) follows from Proposition 2.10 since $q_1^{-1}F \simeq F \boxtimes \mathbf{k}_Y$ and $q_1^!F \simeq F \boxtimes \omega_Y$. To prove (a), denote by \widehat{q}_1 the projection $\widehat{X} \times \widehat{Y} \rightarrow \widehat{X}$. Then $Rq_{1!}H \simeq j_X^{-1}R\widehat{q}_{1!}Rj_{X \times Y!}H$ and similarly $Rq_{1*}H \simeq j_X^{-1}R\widehat{q}_{1*}Rj_{X \times Y*}H$. \square

Remark 2.12. We see in Proposition 2.11 an important difference between constructible sheaves and constructible sheaves up to infinity. Indeed, for usual constructible sheaves, the (proper or non proper) direct image is no more constructible in general.

Corollary 2.13. *Let $f: X_\infty \rightarrow Y_\infty$ be a morphism of b -analytic manifolds and let $F \in D_{\mathbb{R}c}^b(\mathbf{k}_{X_\infty})$ and $G \in D_{\mathbb{R}c}^b(\mathbf{k}_{Y_\infty})$. Then $Rf_*F \simeq D_Y Rf_! D_X F$ and $f^!G \simeq D_X f^{-1} D_Y G$.*

Proof. (i) Both $D_X F$ and $Rf_! D_X F$ are \mathbb{R} -constructible. Then apply [KS90, Exe. VIII.3].
(ii) Similarly, both $D_Y G$ and $f^{-1} D_Y G$ are \mathbb{R} -constructible. Then apply loc. cit. \square

Consider b -analytic manifolds $X_{i\infty} = (X_i, \widehat{X}_i)$, ($i = 1, 2, 3$), and kernels $K_{ij} \in D_{\mathbb{R}c}^b(\mathbf{k}_{X_{ij\infty}})$, $i = 1, 2$, $j = i + 1$. We have already defined in (1.3) the composition of kernels $K_{12} \circ_2 K_{23}$.

Applying Propositions 2.10 and 2.11, we get:

Corollary 2.14. *In the preceding situation, $K_{12} \circ_2 K_{23}$ belongs to $D_{\mathbb{R}c}^b(\mathbf{k}_{X_{13\infty}})$.*

Recall that the composition of kernels is associative (see (1.4)).

2.3 Convolution and γ -topology

In this subsection, we consider a real n -dimensional vector space \mathbb{V} . We consider its projective compactification $\mathbb{P} = (\mathbb{V} \oplus \mathbb{R} \setminus \{0\})/\mathbb{R}^\times$. The pair (\mathbb{V}, \mathbb{P}) is a b-analytic manifold and we set

$$(2.1) \quad \mathbb{V}_\infty = (\mathbb{V}, \mathbb{P}).$$

We simply write \mathbb{V} instead of \mathbb{V}_∞ if there is no risk of confusion.

Convolution

We denote by s the addition map.

$$s: \mathbb{V} \times \mathbb{V} \rightarrow \mathbb{V}, \quad (x, y) \mapsto x + y.$$

Clearly, s is a morphism of b-analytic manifolds.

We define the convolution and the non-proper convolution as follows. For $F, G \in D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{V}_\infty})$, we set

$$F \star G := \mathbf{R}s_!(F \boxtimes G), \quad F \overset{\text{np}}{\star} G := \mathbf{R}s_*(F \boxtimes G).$$

By Propositions 2.11 and 2.10, both $F \star G$ and $F \overset{\text{np}}{\star} G$ belong to $D^b(\mathbb{V}_\infty)$. One checks easily that both convolution operations are commutative and that usual (proper) convolution is associative. Note that, denoting by \mathbb{V}_i ($i = 1, 2$) two copies of \mathbb{V} one has $F_1 \star F_2 \simeq F_1 \boxtimes F_2 \circ_{12} \mathbf{k}_{\Gamma_s}$ where Γ_s is the graph of s in $\mathbb{V}_{12} \times \mathbb{V}$.

Proposition 2.15. *Let $F_i \in D^b(\mathbb{V}_\infty)$, $i = 1, 2, 3$. Then*

$$F_1 \overset{\text{np}}{\star} F_2 \simeq D_{\mathbb{V}}(D_{\mathbb{V}}F_1 \star D_{\mathbb{V}}F_2), \quad (F_1 \overset{\text{np}}{\star} F_2) \overset{\text{np}}{\star} F_3 \simeq F_1 \overset{\text{np}}{\star} (F_2 \overset{\text{np}}{\star} F_3).$$

Proof. (i) The first isomorphism follows from Corollary 2.13.

(ii) follows from (i). □

Remark 2.16. Proposition 2.15 is remarkable since, in general, the operation $\overset{\text{np}}{\star}$ is not associative.

γ -topology

References to the γ -topology and its links with sheaf theory are made to [KS90, KS18]. We consider a real n -dimensional vector space \mathbb{V} . We set $\dot{\mathbb{V}} = \mathbb{V} \setminus \{0\}$ and we recall that \mathbb{V}_∞ is defined in (2.1). Clearly, the antipodal map $a: \mathbb{V} \rightarrow \mathbb{V}$, $x \mapsto -x$, is a morphism of b-analytic manifolds. For a subset A of \mathbb{V} , we denote by A^a its image by the antipodal map.

A subset γ of \mathbb{V} is called a cone if $\mathbb{R}_{>0}\gamma = \gamma$. A closed convex cone γ is proper¹ if $\gamma \cap \gamma^a = \{0\}$.

We consider a cone $\gamma \subset \mathbb{V}$ and we assume:

(2.2) γ is a closed convex proper subanalytic cone with non-empty interior.

¹Of course, the use of ‘‘proper’’ for a cone is not connected to the use of ‘‘proper’’ for a map.

Lemma 2.17. *Let $\gamma \subset \dot{\mathbb{V}}$ be a cone, subanalytic in $\dot{\mathbb{V}}$. Then γ is subanalytic up to infinity.*

Proof. (a) The set γ is subanalytic in \mathbb{V} by [KS90, Prop. 8.3.8 (i)].

(b) Choose a subanalytic norm $\|\cdot\|$ on \mathbb{V} and consider the real analytic isomorphism $f: \dot{\mathbb{V}} \rightarrow \dot{\mathbb{V}}$, $f(x) = x/\|x\|^2$. The map f defines an automorphism of the b-analytic manifold \mathbb{V}_∞ . It is thus enough to check that $f(\gamma)$ is subanalytic in \mathbb{V} . Since this set is a subanalytic cone, this follows from (a). \square

The family of γ -invariant open subsets U of \mathbb{V} (that is, satisfying $U = U + \gamma$) defines a topology, which is called the γ -topology on \mathbb{V} . One denotes by \mathbb{V}_γ the space \mathbb{V} endowed with the γ -topology and one denotes by

$$(2.3) \quad \varphi_\gamma: \mathbb{V} \rightarrow \mathbb{V}_\gamma$$

the continuous map associated with the identity. Note that the closed sets for this topology are the γ^a -invariant closed subsets of \mathbb{V} and that a subset is γ -locally closed if it is the intersection of a γ -closed subset and a γ -open subset.

Lemma 2.18. *Let $A \subset \mathbb{V}$. The conditions below are equivalent:*

- (a) $A = (U + \gamma) \cap \overline{(U + \gamma^a)}$ with U open and subanalytic up to infinity.
- (b) A is the intersection of a γ -closed subset S and a γ -open subset U , both S and U being subanalytic up to infinity.
- (c) A is γ -locally closed and A is subanalytic up to infinity.

Proof. (a) \Rightarrow (b). It remains to check that U being subanalytic up to infinity, $\overline{U + \gamma^a}$ is subanalytic up to infinity. It is enough to check that $U + \gamma^a$ is subanalytic up to infinity. This set is the image of the set $U \times \gamma^a$ by the map $s: \mathbb{V} \times \mathbb{V} \rightarrow \mathbb{V}$, $(x, y) \mapsto x + y$. Hence, the result follows from Proposition 2.6.

(b) \Rightarrow (c) is obvious.

(c) \Rightarrow (a). By [KS18, Prop. 3.4], we may write $A = (U + \gamma) \cap \overline{(U + \gamma^a)}$ with $U = \text{Int}(A)$. Therefore, U is subanalytic up to infinity. \square

Definition 2.19. Let A be a subset of \mathbb{V} . One says that A is *subanalytic γ -locally closed* if A satisfies one of the equivalent conditions in Lemma 2.18.

Let γ be a cone satisfying (2.2). Recall that one denotes by $\gamma^\circ \subset \mathbb{V}^*$ the polar cone.:

$$\gamma^\circ = \{y \in \mathbb{V}^*; \langle x, y \rangle \geq 0 \text{ for all } x \in \gamma\}.$$

γ -constructible sheaves

Consider the full triangulated subcategories of the category $D^b(\mathbf{k}_\mathbb{V})$:

$$(2.4) \quad \begin{cases} D_{\gamma^{oa}}^b(\mathbf{k}_\mathbb{V}) := \{F \in D^b(\mathbf{k}_\mathbb{V}); \text{SS}(F) \subset \mathbb{V} \times \gamma^{oa}\}, \\ D_{\mathbb{R}c, \gamma^{oa}}^b(\mathbf{k}_{\mathbb{V}_\infty}) := D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{V}_\infty}) \cap D_{\gamma^{oa}}^b(\mathbf{k}_\mathbb{V}). \end{cases}$$

We call an object of the category $D_{\mathbb{R}c, \gamma^{oa}}^b(\mathbf{k}_{\mathbb{V}_\infty})$ a γ -constructible sheaf.

Theorem 2.20. *Let $F \in D_{\mathbb{R}c, \gamma^{oa}}^b(\mathbf{k}_{\mathbb{V}\infty})$. Then there exists a finite partition $\mathbb{V} = \bigsqcup_{a \in A} Z_a$ where the Z_a 's are subanalytic γ -locally closed and $F|_{Z_a}$ is constant.*

Proof. This result is proved by Ezra Miller in [Mil20b], using the tools of [Mil20a]. If we make the extra hypothesis that F is PL (piecewise linear) and the cone γ is polyhedral, then this result is proved in [KS18, Th. 3.18]. Note that in loc. cit. the notion of being subanalytic up to infinity is not used and the partition (which is called a stratification there) is only locally finite. However, in our situation, the fact that the partition is finite is implicit in the first part of the proof. \square

Lemma 2.21. *The endofunctor $\mathbf{k}_{\gamma^a}^{\text{np}} \star$ of $D^b(\mathbf{k}_{\mathbb{V}})$ defines a projector $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{V}\infty}) \rightarrow D_{\mathbb{R}c, \gamma^{oa}}^b(\mathbf{k}_{\mathbb{V}\infty})$.*

Proof. We know by [KS90, Prop. 5.2.3] that the functor $\varphi_{\gamma}^{-1}R\varphi_{\gamma*}: D^b(\mathbf{k}_{\mathbb{V}}) \rightarrow D_{\gamma^{oa}}^b(\mathbf{k}_{\mathbb{V}})$ is a projector and we know by [KS90, Prop. 3.5.4] that the two functors $\varphi_{\gamma}^{-1}R\varphi_{\gamma*}$ and $\mathbf{k}_{\gamma^a}^{\text{np}} \star$ are isomorphic. Moreover, the functor $\mathbf{k}_{\gamma^a}^{\text{np}} \star$ sends $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{V}\infty})$ to itself by Proposition 2.11. \square

Remark 2.22. In general, non proper convolution is not defined on $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{V}})$ and, in particular, even if γ is subanalytic, the functor $\mathbf{k}_{\gamma^a}^{\text{np}} \star$ does not send $D_{\mathbb{R}c}^b(\mathbf{k}_{\mathbb{V}})$ to itself.

3 A short review on constructible functions

From now on and until the end of this paper, we assume that \mathbf{k} is a field of *characteristic zero*.

In this section, we recall without proofs the main constructions and results on constructible functions. References are made to [Sch91] and [KS90, § 9.7].

3.1 From constructible sheaves to constructible functions

Definition 3.1. A function $\varphi: X \rightarrow \mathbb{Z}$ is constructible if:

- (i) for all $m \in \mathbb{Z}$, $\varphi^{-1}(m)$ is subanalytic in X ,
- (ii) the family $\{\varphi^{-1}(m)\}_{m \in \mathbb{Z}}$ is locally finite.

Notation 3.2. For a locally closed subanalytic subset $S \subset X$, we denote by $\mathbf{1}_S$ the characteristic function of S (with values 1 on S and 0 elsewhere). For $a \in X$ we also set $\delta_a = \mathbf{1}_{\{a\}}$.

The next result is well known. Note that the implication (b) \Rightarrow (d) follows from the triangulation theorem for compact subanalytic subsets.

Lemma 3.3. *Let φ be a \mathbb{Z} -valued function on X . The conditions below are equivalent.*

- (a) φ is constructible,

- (b) *there exist a locally finite family of subanalytic locally closed subsets $\{Z_i\}_{i \in I}$ and $c_i \in \mathbb{Z}$ such that $\varphi = \sum_i c_i \mathbf{1}_{Z_i}$,*
- (c) *there exist a subanalytic stratification $\{Z_i\}_{i \in I}$ and $c_i \in \mathbb{Z}$ such that $\varphi = \sum_i c_i \mathbf{1}_{Z_i}$,*
- (d) *same as (b) assuming moreover each Z_i compact and contractible.*

Notation 3.4. One denotes by $\mathcal{CF}(X)$ the group of constructible functions on X and by \mathcal{CF}_X the presheaf $U \mapsto \mathcal{CF}(U)$.

Proposition 3.5. *The presheaf \mathcal{CF}_X is a sheaf on X .*

Proof. (i) Clearly, the presheaf $U \mapsto \mathcal{CF}(U)$ is separated.

(ii) Let $X = \bigcup_{a \in A} U_a$ be an open covering of X and let φ be a \mathbb{Z} -valued function on X such that $\varphi|_{U_a}$ is constructible on U_a . For $m \in \mathbb{Z}$, set $Z_m := \varphi^{-1}(m)$ and $Z_{m,a} = Z_m \cap U_a$. Each $Z_{m,a}$ is subanalytic in U_a , which implies that Z_m is subanalytic in X . Moreover, the family $\{Z_{m,a}\}_m$ being locally finite in U_a , the family $\{Z_m\}_m$ is locally finite in X . Hence, φ is constructible on X . The same argument holds when replacing X with an open subset $U \subset X$. \square

Recall an important theorem (see [KS90, Th. 9.7.1]) which clarifies the notion of constructible function. First, denote by χ_{loc} the local Euler-Poincaré index:

$$(3.1) \quad \chi_{\text{loc}}: \text{Ob}(\mathbb{D}_{\mathbb{R}c}^b(\mathbf{k}_X)) \rightarrow \mathcal{CF}(X), \quad \chi_{\text{loc}}(F)(x) = \sum_i (-1)^i \dim H^i(F_x).$$

Denote by $\mathbb{K}(\mathcal{C})$ the Grothendieck group of either an abelian or a triangulated category \mathcal{C} , and recall that if \mathcal{C} is abelian then $\mathbb{K}(\mathcal{C}) \xrightarrow{\simeq} \mathbb{K}(\mathbb{D}^b(\mathcal{C}))$. Recall that by its construction, any additive map $\text{Ob}(\mathcal{C}) \rightarrow \mathbb{Z}$ factorizes uniquely through $\mathbb{K}(\mathcal{C})$. Also recall that if $F: \mathcal{C} \rightarrow \mathcal{C}'$ is a triangulated functor, then it defines a linear map $\mathbb{K}(\mathcal{C}) \rightarrow \mathbb{K}(\mathcal{C}')$.

In the sequel, we set for short

$$\mathbb{K}_{\mathbb{R}c}(\mathbf{k}_X) := \mathbb{K}(\mathbb{D}_{\mathbb{R}c}^b(\mathbf{k}_X)).$$

The tensor product on $\mathbb{D}_{\mathbb{R}c}^b(\mathbf{k}_X)$ defines a ring structure on $\mathbb{K}_{\mathbb{R}c}(\mathbf{k}_X)$, with unit the image of the constant sheaf \mathbf{k}_X .

Theorem 3.6. *Let X be a real analytic manifold. Then the map χ_{loc} defines an isomorphism of commutative unital algebras (we keep the same notation) $\chi_{\text{loc}}: \mathbb{K}_{\mathbb{R}c}(\mathbf{k}_X) \xrightarrow{\simeq} \mathcal{CF}(X)$.*

Note that if $\chi_{\text{loc}}(F) = \varphi$ and $S := \text{supp}(\varphi)$, then $\chi_{\text{loc}}(F_S) = \varphi$. Hence, given $\varphi \in \mathcal{CF}(X)$, we may always represent φ with a constructible sheaf of same support. We have the general “principle” that we shall make explicit in the sequel:

The operations on constructible functions are the image by the local Euler-Poincaré index χ_{loc} of the corresponding operations on constructible sheaves.

In the sequel, we shall also encounter the global Euler-Poincaré indices of a sheaf F (assuming that these indices are finite):

$$(3.2) \quad \chi(F) = \chi(\text{R}\Gamma(X; F)), \quad \chi_c(F) = \chi(\text{R}\Gamma_c(X; F)).$$

For a locally closed subanalytic subset Z of X , we also set

$$(3.3) \quad \chi(Z) = \chi(\mathbf{k}_Z) = \chi(\mathrm{R}\Gamma(Z; \mathbf{k}_Z)), \quad \chi_c(Z) = \chi_c(\mathbf{k}_Z) = \chi(\mathrm{R}\Gamma_c(Z; \mathbf{k}_Z)).$$

Recall that denoting by $j: Z \hookrightarrow X$ the embedding, $\mathbf{k}_{XZ} = j_!\mathbf{k}_Z$. Hence, $\mathrm{R}\Gamma_c(Z; \mathbf{k}_Z) \simeq \mathrm{R}\Gamma_c(X; \mathbf{k}_{XZ})$ but $\mathrm{R}\Gamma(Z; \mathbf{k}_Z) \simeq \mathrm{R}\Gamma(X; \mathrm{R}j_*\mathbf{k}_Z) \neq \mathrm{R}\Gamma(X; \mathbf{k}_{XZ})$ in general.

3.2 Operations

Internal operations

The sum on $\mathcal{CF}(X)$ is the image by χ_{loc} of the direct sum for sheaves, the unit $\mathbf{1}_X$ is the image of the constant sheaf \mathbf{k}_X , the map $\varphi \mapsto -\varphi$ corresponds to the shift $F \mapsto F[+1]$ and the usual product on $\mathcal{CF}(X)$ is the image of the tensor product.

External product

For two real analytic manifolds X and Y , one defines the morphism

$$(3.4) \quad \boxtimes: \mathcal{CF}_X \boxtimes \mathcal{CF}_Y \rightarrow \mathcal{CF}_{X \times Y}, \quad (\varphi \boxtimes \psi)(x, y) = \varphi(x)\psi(y).$$

Inverse image or composition

Let $f: X \rightarrow Y$ be a morphism of real analytic manifolds. One defines the inverse image morphism

$$(3.5) \quad f^*: f^{-1}\mathcal{CF}_Y \rightarrow \mathcal{CF}_X, \quad (f^*\psi)(x) = \psi(f(x)) \text{ for } \psi \in \mathcal{CF}(Y).$$

(Recall that a morphism $f^{-1}\mathcal{CF}_Y \rightarrow \mathcal{CF}_X$ is nothing but a morphism $\mathcal{CF}_Y \rightarrow f_*\mathcal{CF}_X$.)

Inverse images are functorial, that is, if $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ are morphisms of manifolds, then:

$$f^* \circ g^* = (g \circ f)^*.$$

Direct image or integral

Recall first that, if K is a subanalytic compact subset of X , then the Euler-Poincaré index $\chi(K)$ is defined by $\chi(K) = \sum_j (-1)^j b_j(K)$ where $b_j(K) = \dim_{\mathbb{Q}} H^j(K; \mathbb{Q}_K) = \dim_{\mathbf{k}} H^j(K; \mathbf{k}_K)$ (recall that \mathbf{k} is a field of characteristic 0 and recall (3.2)). In particular, if K is contractible, then $\chi(K) = 1$. One sets

$$(3.6) \quad \int_X \mathbf{1}_K = \chi(K).$$

If φ has compact support, one may assume that the sum in Lemma 3.3 (d) is finite, and one checks (one can use either Theorem 3.6 or the triangulation theorem for subanalytic sets) that the integer $\sum_i c_i$ depends only on φ , not on its decomposition. One sets:

$$\int_X \varphi = \sum_i c_i.$$

In particular, if Z is locally closed relatively compact and subanalytic in X , then (see (3.3)):

$$(3.7) \quad \int_X \mathbf{1}_Z = \chi_c(Z).$$

One shall be aware that the integral is not positive, that is

$$\varphi \geq 0 \text{ does not imply } \int_X \varphi \geq 0.$$

For example, take $X = \mathbb{R}$ and $\varphi = \mathbf{1}_{(-1,1)}$. Hence, $\varphi \geq 0$ and $\int_{\mathbb{R}} \varphi = -1$.

Let $f: X \rightarrow Y$ be a morphism of real analytic manifolds. One defines the direct image morphism

$$(3.8) \quad \int_X : f_! \mathcal{CF}_X \rightarrow \mathcal{CF}_Y, \quad \left(\int_f \varphi \right)(y) = \int_X \mathbf{1}_{f^{-1}(y)} \cdot \varphi.$$

Recall that a section of $f_! \mathcal{CF}_Y$ on an open subset $U \subset X$ is a section of $\mathcal{CF}_X(f^{-1}U)$ such that f is proper on its support. Hence the integral makes sense as a function but it is not obvious that it is a constructible function. This follows for example from the corresponding result for direct images of constructible sheaves. Indeed, let $F \in D_{\mathbb{R}c}^b(\mathbf{k}_X)$ be such that $\chi_{\text{loc}}(F) = \varphi$ and $\text{supp}(F) = \text{supp}(\varphi)$. Then $\int_f \varphi = \chi_{\text{loc}}(\mathbf{R}f_! F)$.

Direct images are functorial, that is, if $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ are morphisms of manifolds, then:

$$\int_g \circ \int_f = \int_{g \circ f}.$$

Duality

On X , the dual of a constructible function is the image by χ_{loc} of the duality functor D_X for sheaves. For $F \in D^b(\mathbf{k}_X)$ and $x_0 \in X$, one has

$$(D_X F)_{x_0} \simeq (\mathbf{R}\Gamma_{x_0}(F))^*,$$

where $*$ denotes the duality functor for \mathbf{k} -vector spaces. Since F is constructible, there exists a local chart and $\varepsilon_0 > 0$ such that, denoting by $B_\varepsilon(x_0)$ the open ball with center x_0 and radius $\varepsilon > 0$ in this chart, one has for $0 < \varepsilon \leq \varepsilon_0$:

$$\mathbf{R}\Gamma_{x_0}(F) \simeq \mathbf{R}\Gamma_c(B_\varepsilon(x_0); F) \simeq \mathbf{R}a_{X!}(F \otimes \mathbf{k}_{B_\varepsilon(x_0)}).$$

Hence, one defines the dual of a constructible function φ on X as follows. Let $x_0 \in X$, and choose a local chart in a neighborhood of x_0 and $\varepsilon > 0$ as above. One sets

$$(3.9) \quad (D_X \varphi)(x_0) = \int_X \varphi \cdot \mathbf{1}_{B_\varepsilon(x_0)}.$$

The integral $\int_X \varphi \cdot \mathbf{1}_{B_\varepsilon(x_0)}$ neither depends on the local chart nor on ε , for $0 < \varepsilon \leq \varepsilon_0$, for some $\varepsilon_0 > 0$ depending on x_0 .

We get a morphism of sheaves $D_X: \mathcal{CF}_X \rightarrow \mathcal{CF}_X$ and this morphism is an involution, that is,

$$D_X \circ D_X \simeq \text{id}_X.$$

Moreover, duality commutes with integration. Assuming that f is proper on the support of φ , one has:

$$(3.10) \quad D_Y\left(\int_f \varphi\right) = \int_f D_X(\varphi).$$

By mimicking a classical formula for constructible sheaves, one sets

$$(3.11) \quad \text{hom}(\varphi, \psi) := D_X(D_X\psi \cdot \varphi).$$

Example 3.7. Let Z be a closed subanalytic subset of X and assume that Z is a C^0 -manifold of dimension d with boundary ∂Z . Set $A = Z \setminus \partial Z$. Hence, locally on X , $Z \subset X$ is topologically isomorphic to $\overline{U} \subset \mathbb{R}^n$ where U is a convex open subset of $\mathbb{R}^d \subset \mathbb{R}^n$ and $A \simeq U$. We thus have

$$(3.12) \quad D_X \mathbf{1}_Z = (-1)^d \mathbf{1}_A$$

Moreover

$$\int_X \mathbf{1}_{\partial Z} = \int_X \mathbf{1}_Z - \int_X \mathbf{1}_A = (1 - (-1)^d) \int_X \mathbf{1}_Z.$$

When Z is a closed convex polyhedron, one recovers the classical Euler formula.

Other operations

In fact, most (if not all) operations on constructible sheaves admit a counterpart in the language of constructible functions. In [KS90, Def. 9.7.8] one defines the specialization ν_M along a submanifold M , its Fourier-Sato transform, the microlocalization μ_M and μhom :

$$\begin{aligned} \nu_M: \mathcal{CF}(X) &\rightarrow \mathcal{CF}_{\mathbb{R}^+}(T_M X), & \mu_M: \mathcal{CF}(X) &\rightarrow \mathcal{CF}_{\mathbb{R}^+}(T_M^* X) \\ \mu\text{hom}: \mathcal{CF}(X) \times \mathcal{CF}(X) &\rightarrow \mathcal{CF}_{\mathbb{R}^+}(T^* X). \end{aligned}$$

One can also define the micro-support of $\varphi \in \mathcal{CF}(X)$ by setting

$$(3.13) \quad \text{SS}(\varphi) = \text{supp}(\mu\text{hom}(\varphi, \varphi)).$$

4 Constructible functions up to infinity

4.1 Definition

Definition 4.1. Let $X_\infty = (X, \widehat{X})$ be a b-analytic manifold. A function $\varphi: X \rightarrow \mathbb{Z}$ is constructible up to infinity if:

- (i) for all $m \in \mathbb{Z}$, $\varphi^{-1}(m)$ is subanalytic up to infinity,
- (ii) the family $\{\varphi^{-1}(m)\}_{m \in \mathbb{Z}}$ is finite.

We denote by $\mathcal{CF}(X_\infty)$ the space of constructible functions up to infinity.

For any function φ on X , we denote by $j_{X!}\varphi$ the function on \widehat{X} obtained as the function φ on X extended by 0 on $\widehat{X} \setminus X$.

Lemma 4.2. *Let $\varphi \in \mathcal{CF}(X)$. The conditions below are equivalent.*

- (a) *The function φ is constructible up to infinity,*
- (b) *The function $j_{X!}\varphi$ belongs to $\mathcal{CF}(\widehat{X})$.*
- (c) *There exists $\psi \in \mathcal{CF}(\widehat{X})$ such that $\varphi = \psi|_X$.*
- (d) *There exist a finite family of locally closed subsets subanalytic up to infinity $\{Z_i\}_{i \in I}$ and $c_i \in \mathbb{Z}$ such that $\varphi = \sum_i c_i \mathbf{1}_{Z_i}$.*

Proof. (a) \Rightarrow (b). By the hypothesis, one may write $\varphi = \sum_i c_i \mathbf{1}_{Z_i}$ where the sum is finite and the Z_i 's are subanalytic up to infinity. Therefore, $\mathbf{1}_{Z_i} \in \mathcal{CF}(\widehat{X})$ and the result follows from Lemma 3.3.

(b) \Rightarrow (c) is obvious.

(c) \Rightarrow (d) and (c) \Rightarrow (a). By definition, for each $m \in \mathbb{Z}$, $Z_m := \psi^{-1}(m)$ is subanalytic in \widehat{X} and the family $\{Z_m\}_m$ is locally finite. Therefore, $Z_m \cap X$ is subanalytic in X and X being relatively compact, the family $\{X \cap Z_m\}_m$ is finite.

(d) \Rightarrow (b) is obvious. □

Clearly, $\mathcal{CF}(X_\infty)$ is a subalgebra of $\mathcal{CF}(X)$.

Let us denote by \mathcal{CF}_{X_∞} the presheaf on $X_{\infty\text{sa}}$ given by $U \mapsto \mathcal{CF}(U_\infty)$.

Proposition 4.3. *The presheaf \mathcal{CF}_{X_∞} is a sheaf on $X_{\infty\text{sa}}$.*

The proof is straightforward.

Recall Theorem 3.6 and denote now by $\mathbb{K}_{\mathbb{R}c}(\mathbf{k}_{X_\infty})$ the Grothendieck group of the category $D_{\mathbb{R}c}^b(\mathbf{k}_{X_\infty})$.

Theorem 4.4. *The isomorphism of commutative unital algebras $\chi_{\text{loc}}: \mathbb{K}_{\mathbb{R}c}(\mathbf{k}_X) \xrightarrow{\sim} \mathcal{CF}(X)$ induces an isomorphism $\chi_{\text{loc}}: \mathbb{K}_{\mathbb{R}c}(\mathbf{k}_{X_\infty}) \xrightarrow{\sim} \mathcal{CF}(X_\infty)$.*

Proof. (i) The map χ_{loc} takes its values in $\mathcal{CF}(X_\infty)$. Indeed, for $F \in D_{\mathbb{R}c}^b(\mathbf{k}_{X_\infty})$, $\chi_{\text{loc}}(F) = j_X^*(\chi_{\text{loc}}(j_{X!}F))$.

(ii) The map $\chi_{\text{loc}}: \mathbb{K}_{\mathbb{R}c}(\mathbf{k}_{X_\infty}) \rightarrow \mathcal{CF}(X_\infty)$ is injective by the same arguments as in the proof of [KS90, Th. 9.7.1].

(iii) The map χ_{loc} is surjective since for Z locally closed and subanalytic up to infinity, $\mathbf{1}_Z = \chi_{\text{loc}}(\mathbf{k}_Z)$ and \mathbf{k}_Z is constructible up to infinity. □

4.2 Operations

Lemma 4.5. *If $\varphi \in \mathcal{CF}(X_\infty)$, then $D_X\varphi \in \mathcal{CF}(X_\infty)$.*

Proof. The result follows from Lemma 4.2 (c) since duality commutes with restriction to an open subset. \square

Let $\varphi \in \mathcal{CF}(X_\infty)$. One sets

$$(4.1) \quad j_{X*}\varphi = D_{\widehat{X}}j_{X!}D_X\varphi.$$

The next result follows from the corresponding result for sheaves.

Lemma 4.6. *If $\varphi \in \mathcal{CF}(X_\infty)$ has compact support in X , then $j_{X*}\varphi = j_{X!}\varphi$.*

Proposition 4.7. *Let X_∞ and Y_∞ be two b-analytic manifolds.*

- (a) *Let $\varphi \in \mathcal{CF}(X_\infty)$ and $\psi \in \mathcal{CF}(Y_\infty)$. Then the function $\varphi \boxtimes \psi$, defined by $(\varphi \boxtimes \psi)(x, y) = \varphi(x)\psi(y)$, belongs to $\mathcal{CF}((X \times Y)_\infty)$.*
- (b) *Let $f: X_\infty \rightarrow Y_\infty$ be a morphism of b-analytic manifolds and let $\psi \in \mathcal{CF}(Y_\infty)$. Then the function $f^*\psi$ defined by $f^*\psi(x) = \psi(f(x))$ belongs to $\mathcal{CF}(X_\infty)$.*

In other words we have extended the morphisms (3.5) and (3.4) to b-analytic manifolds.

Proof. (a) Apply Lemma 4.2.

(b) Apply Proposition 2.6 together with Definition 4.1. \square

Although we shall not use it, let us mention that one can also define the internal hom and the exceptional inverse image by the formulas

$$(4.2) \quad \begin{aligned} \text{hom}(\varphi, \psi) &:= D_X(D_X\psi \cdot \varphi), \quad \varphi, \psi \in \mathcal{CF}(X_\infty), \\ f^!\psi &:= D_X f^*(D_Y\psi), \quad \psi \in \mathcal{CF}(Y_\infty). \end{aligned}$$

Now we study the integrals of constructible functions up to infinity. One can define two integrals of $\varphi \in \mathcal{CF}(X_\infty)$. One sets

$$(4.3) \quad \int_X \varphi := \int_{\widehat{X}} j_{X!}\varphi, \quad \int_X^{\text{np}} \varphi := \int_X D_X\varphi.$$

Recall notations (3.3).

Lemma 4.8. (a) *One has $\int_X^{\text{np}} \varphi = \int_{\widehat{X}} j_{X*}\varphi$.*

(b) *Let Z be a locally closed subset of X subanalytic up to infinity. Then $\int_X \mathbf{1}_Z = \chi_c(Z)$ and $\int_X^{\text{np}} \mathbf{1}_Z = \chi(Z)$.*

(c) *The integrals $\int_X \varphi$ and $\int_X^{\text{np}} \varphi$ do not depend on the choice of \widehat{X} .*

Proof. (a) follows from

$$\int_{\widehat{X}} j_{X*} \varphi = \int_{\widehat{X}} D_{\widehat{X}} j_{X!} D_X \varphi = \int_{\widehat{X}} j_{X!} D_X \varphi$$

where the last equality follows from (3.10) applied with $Y = \text{pt}$.

(b) Recall (3.7). Also recall that a_Z is the map $Z \rightarrow \text{pt}$ and similarly with $a_{\widehat{X}}$. Denoting by j_Z the embedding $Z \hookrightarrow \widehat{X}$, we have

$$\begin{aligned} \int_X \mathbf{1}_Z &= \chi(\text{Ra}_{\widehat{X}!} \text{R}j_{Z!} \mathbf{k}_Z) = \chi(\text{Ra}_{Z!} \mathbf{k}_Z) = \chi_c(Z), \\ \int_X^{\text{np}} \mathbf{1}_Z &= \chi(\text{Ra}_{\widehat{X}*} \text{R}j_{Z*} \mathbf{k}_Z) = \chi(\text{Ra}_{Z*} \mathbf{k}_Z) = \chi(Z). \end{aligned}$$

(We have used the fact that $\text{R}j_{Z*} \mathbf{k}_Z$ has compact support in \widehat{X} .)

(c) follows from (b). □

Example 4.9. Let $X = \mathbb{R}$. Then:

- (i) One has $\int_{\mathbb{R}} \mathbf{1}_{\mathbb{R}} = -1$, $\int_{\mathbb{R}}^{\text{np}} \mathbf{1}_{\mathbb{R}} = 1$.
- (ii) Let $U = (-\infty, b)$ with $-\infty < b < \infty$. Then $\int_{\mathbb{R}} \mathbf{1}_U = -1$, $\int_{\mathbb{R}}^{\text{np}} \mathbf{1}_U = 0$.
- (iii) Let $Z = (-\infty, b]$ with $-\infty < b < +\infty$. Then $\int_{\mathbb{R}} \mathbf{1}_Z = 0$, $\int_{\mathbb{R}}^{\text{np}} \mathbf{1}_Z = 1$.
- (iv) Let $S = [a, b]$ with $-\infty < a \leq b < +\infty$. Then $\int_{\mathbb{R}} \mathbf{1}_S = \int_{\mathbb{R}}^{\text{np}} \mathbf{1}_S = 1$.
- (v) Let $Z = [a, b)$ with $-\infty < a \leq b < +\infty$. Then $\int_{\mathbb{R}} \mathbf{1}_Z = \int_{\mathbb{R}}^{\text{np}} \mathbf{1}_Z = 0$.

Indeed, (i) is obvious. Let U be as in (ii). Then U is topologically isomorphic to \mathbb{R} and we get $\int_{\mathbb{R}} \mathbf{1}_U = -1$. By the additivity of the integral, we deduce that for Z as in (iii), $\int_{\mathbb{R}} \mathbf{1}_Z = 0$. By Lemma 4.8, we get $\int_{\mathbb{R}}^{\text{np}} \mathbf{1}_U = 0$ and by additivity, $\int_{\mathbb{R}}^{\text{np}} \mathbf{1}_Z = 1$. Finally, (iv) and (v) are obvious.

Let $f: X_{\infty} \rightarrow Y_{\infty}$ be a morphism of b-analytic manifolds and let $\varphi \in \mathcal{CF}(X_{\infty})$. Similarly as in (3.8), one sets for $y \in Y$:

$$(4.4) \quad \left(\int_f \varphi \right)(y) = \int_X \mathbf{1}_{f^{-1}(y)} \cdot \varphi.$$

Of course, when $Y = \text{pt}$, one recovers (4.3).

Lemma 4.10. *The function $\int_f \varphi$ defined by (4.4) belongs to $\mathcal{CF}(Y_{\infty})$.*

Proof. Let us choose $F \in D^b(\mathbf{k}_{X_{\infty}})$ such that $\chi_{\text{loc}}(F) = \varphi$. Then $(\int_f \varphi)(y) = \chi_{\text{loc}}(\text{R}f_! F)$ and $\text{R}f_! F \in D^b(\mathbf{k}_{Y_{\infty}})$. □

Hence, we have constructed a morphism

$$\int_f: \mathcal{CF}_{X_{\infty}} \rightarrow \mathcal{CF}_{Y_{\infty}}, \quad \varphi \mapsto \int_f \varphi.$$

We also define

$$\int_f^{\text{np}} : \mathcal{CF}_{X_\infty} \rightarrow \mathcal{CF}_{Y_\infty}, \quad \int_f^{\text{np}} \varphi := D_Y \int_f D_X \varphi.$$

The next results are easily checked.

- If f is proper on $\text{supp}(\varphi)$, then $\int_f \varphi = \int_f^{\text{np}} \varphi$.
- If $\varphi = \chi_{\text{loc}}(F)$ for some $F \in D_{\mathbb{R}c}^b(\mathbf{k}_{X_\infty})$, then $\int_f \varphi = \chi_{\text{loc}}(\mathbf{R}f_! F)$ and $\int_f^{\text{np}} \varphi = \chi_{\text{loc}}(\mathbf{R}f_* F)$.
- Let $g: Y_\infty \rightarrow Z_\infty$ be another morphism of b-analytic manifolds. Then

$$\int_{g \circ f} \varphi = \int_g \int_f \varphi, \quad \int_{g \circ f}^{\text{np}} \varphi = \int_g^{\text{np}} \int_f^{\text{np}} \varphi.$$

Base change formula

Consider two morphisms $f: X_\infty \rightarrow Z_\infty$ and $g: Y_\infty \rightarrow Z_\infty$ of b-analytic manifolds and consider ² a Cartesian square of topological spaces

$$(4.5) \quad \begin{array}{ccc} W & \xrightarrow{f'} & Y \\ g' \downarrow & & \downarrow g \\ X & \xrightarrow{f} & Z. \end{array}$$

Recall that the square is Cartesian means that W is isomorphic to the space $\{(x, y) \in X \times Y; f(x) = g(y)\}$. We consider W as a closed subanalytic subset of $X \times Y$.

Proposition 4.11. *Consider the square (4.5) and let $\varphi \in \mathcal{CF}(X_\infty)$. Then $\int_{f'}(g'^* \varphi)$ is well defined, belongs to $\mathcal{CF}(Y_\infty)$ and*

$$(4.6) \quad g^* \int_f \varphi = \int_{f'}(g'^* \varphi).$$

Proof. Choose $F \in D_{\mathbb{R}c}^b(\mathbf{k}_{\widehat{X}})$ such that $\chi_{\text{loc}}(F) = \varphi$. Then apply the base change formula for sheaves. \square

The projection formula

Proposition 4.12. *Let $f: X_\infty \rightarrow Y_\infty$ be a morphism of b-analytic manifolds, let $\varphi \in \mathcal{CF}(X_\infty)$ and $\psi \in \mathcal{CF}(Y_\infty)$. Then*

$$(4.7) \quad \int_f (\varphi \cdot f^* \psi) = \psi \int_f \varphi.$$

²In [Sch20] it was made reference to the notion of a Cartesian square in the category of b-analytic manifolds, a notion which should have been defined more precisely and that we avoid here.

Proof. Choose $F \in D_{\mathbb{R}c}^b(\mathbf{k}_{\widehat{X}})$ such that $\chi_{\text{loc}}(F) = \varphi$ and choose $G \in D_{\mathbb{R}c}^b(\mathbf{k}_{\widehat{Y}})$ such that $\chi_{\text{loc}}(G) = \psi$. Then apply the projection formula for sheaves. \square

Example 4.13. Equality (4.7) is no longer true when replacing \int_f with \int_f^{np} . Set $X = \mathbb{R}^2$ with coordinates (y, t) and $Y = \mathbb{R}$, f being the first projection. Let $\varphi = \mathbf{1}_S$ with $S = \{(y, t); t = 1/(1 - y^2), -1 < y < 1\}$ and let $\psi = \mathbf{1}_Z$ with $Z = (-1, 1)$. One checks easily that φ is subanalytic up to infinity when choosing for example for \widehat{X} the projective compactification of \mathbb{R}^2 . We have $\mathbf{1}_S \cdot f^* \mathbf{1}_Z = \mathbf{1}_S$, $\int_f \mathbf{1}_S = \mathbf{1}_Z$ and $D_X \mathbf{1}_S = -\mathbf{1}_S$ (see Example 3.7). Hence,

$$\begin{aligned} \int_f^{\text{np}} \mathbf{1}_S \cdot f^* \mathbf{1}_Z &= \int_f^{\text{np}} \mathbf{1}_S = D_Y \int_f D_X \mathbf{1}_S = -D_Y \mathbf{1}_Z = \mathbf{1}_{[-1, 1]}, \\ \mathbf{1}_Z \cdot \int_f^{\text{np}} \mathbf{1}_S &= \mathbf{1}_Z \cdot \mathbf{1}_{[-1, 1]} = \mathbf{1}_{(-1, 1)}. \end{aligned}$$

Composition of kernels

Recall Diagram 1.1 when replacing the manifolds X_i with b-analytic manifolds $X_{i\infty}$ ($i = 1, 2, 3$). Let $\lambda_{12} \in \mathcal{CF}(X_{12\infty})$ and $\lambda_{23} \in \mathcal{CF}(X_{23\infty})$. It follows from Proposition 4.7 that the function

$$(4.8) \quad \lambda_{12} \circ_2 \lambda_{23} := \int_{q_{13}} q_{12}^* \lambda_{12} \cdot q_{23}^* \lambda_{23}.$$

is well-defined and belongs to $\mathcal{CF}(X_{13\infty})$. Moreover

Theorem 4.14. *Let $\lambda_{ij} \in \mathcal{CF}(X_{ij\infty})$ ($i = 1, 2, 3, 4$, $j = i + 1$). One has*

$$(\lambda_{12} \circ_2 \lambda_{23}) \circ_3 \lambda_{34} = \lambda_{12} \circ_2 (\lambda_{23} \circ_3 \lambda_{34}) \in \mathcal{CF}(X_{14\infty}).$$

One can prove this theorem by mimicking the classical proof for sheaves, using now Propositions 4.11 and 4.12. One can also prove this result by replacing each λ_{ij} with a kernel $K_{ij} \in D_{\mathbb{R}c}^b(\mathbf{k}_{X_{ij\infty}})$.

4.3 γ -constructible functions

Let \mathbb{V} and \mathbb{V}_∞ be as in § 2.3. We define the convolution and the non-proper convolution similarly as for sheaves (see Proposition 2.15). For $\varphi, \psi \in \mathcal{CF}(\mathbb{V}_\infty)$, we set

$$\varphi \star \psi := \int_s \varphi \boxtimes \psi, \quad \varphi \overset{\text{np}}{\star} \psi := \int_s^{\text{np}} \varphi \boxtimes \psi.$$

By the preceding results, both $\varphi \star \psi$ and $\varphi \overset{\text{np}}{\star} \psi$ belong to $\mathcal{CF}(\mathbb{V}_\infty)$. Note that

$$\varphi \star \psi = \psi \star \varphi, \quad \varphi \overset{\text{np}}{\star} \psi = \psi \overset{\text{np}}{\star} \varphi.$$

Lemma 4.15. *Let $\varphi_i \in \mathcal{CF}(\mathbb{V}_\infty)$, $i = 1, 2, 3$. Then*

$$\varphi_1 \overset{\text{np}}{\star} \varphi_2 = D_X(D_X \varphi_1 \star D_X \varphi_2), \quad (\varphi_1 \overset{\text{np}}{\star} \varphi_2) \overset{\text{np}}{\star} \varphi_3 = \varphi_1 \overset{\text{np}}{\star} (\varphi_2 \overset{\text{np}}{\star} \varphi_3).$$

Proof. The first equality follows from the definition of \int^{np} (see (4.4)) and the second equality follows from the first one. \square

Definition 4.16. Let $\varphi \in \mathcal{CF}(\mathbb{V}_\infty)$. We say that φ is γ -constructible if there exists a finite covering $\mathbb{V} = \bigcup_a Z_a$ such that $\varphi = \sum_a c_a \mathbf{1}_{Z_a}$ and the Z_a 's are subanalytic γ -locally closed subsets of \mathbb{V} . We denote by $\mathcal{CF}(\mathbb{V}_\gamma)$ the space of γ -constructible functions on \mathbb{V} .

By construction, we have $\mathcal{CF}(\mathbb{V}_\gamma) \subset \mathcal{CF}(\mathbb{V}_\infty)$.

We denote by $\mathbb{K}_{\mathbb{R}c, \gamma^{\text{oa}}}(\mathbf{k}_{\mathbb{V}_\infty})$ the Grothendieck group of $D_{\mathbb{R}c, \gamma^{\text{oa}}}^b(\mathbf{k}_{\mathbb{V}_\infty})$.

Theorem 4.17. *The isomorphism of commutative unital algebras $\chi_{\text{loc}}: \mathbb{K}_{\mathbb{R}c}(\mathbf{k}_{\mathbb{V}}) \xrightarrow{\sim} \mathcal{CF}(\mathbb{V})$ induces an isomorphism $\chi_{\text{loc}}: \mathbb{K}_{\mathbb{R}c, \gamma^{\text{oa}}}(\mathbf{k}_{X_\infty}) \xrightarrow{\sim} \mathcal{CF}(\mathbb{V}_\gamma)$.*

Proof. (i) It follows from Theorem 2.20 that the map χ_{loc} takes its values in $\mathcal{CF}(\mathbb{V}_\gamma)$.
(ii) The map χ_{loc} is injective by Lemma 2.21. Indeed, if \mathcal{A} is a full triangulated category of a triangulated category \mathcal{T} and if there is a projector $P: \mathcal{T} \rightarrow \mathcal{A}$, then P induces a projector $\mathbb{K}(P): \mathbb{K}(\mathcal{T}) \rightarrow \mathbb{K}(\mathcal{A})$. In particular, $\mathbb{K}(\mathcal{A})$ is a subgroup of $\mathbb{K}(\mathcal{T})$.
(iii) The map χ_{loc} is surjective since for Z subanalytic γ -locally closed, $\mathbf{1}_Z = \chi_{\text{loc}}(\mathbf{k}_Z)$ and $\mathbf{k}_Z \in D_{\mathbb{R}c}^b(\mathbb{V}_\infty)$. Moreover, $\text{SS}(\mathbf{k}_Z) \subset \mathbb{V} \times \gamma^{\text{oa}}$ by [KS18, Cor. 1.8]. \square

The projector of Lemma 2.21 allows us to construct a projector $\mathcal{CF}(\mathbb{V}_\infty) \rightarrow \mathcal{CF}(\mathbb{V}_\gamma)$.

Proposition 4.18. (a) *Let $\varphi \in \mathcal{CF}(\mathbb{V}_\infty)$. Then $\varphi \star^{\text{np}} \mathbf{1}_{\gamma^{\text{oa}}}$ belongs to $\mathcal{CF}(\mathbb{V}_\gamma)$.*

(b) *If $\varphi \in \mathcal{CF}(\mathbb{V}_\gamma)$, then $\varphi \star^{\text{np}} \mathbf{1}_{\gamma^{\text{oa}}} = \varphi$.*

Proof. The result follows from Theorem 4.17, Lemma 2.21 and the fact that the operation \star^{np} commutes with χ_{loc} . \square

5 Correspondences for constructible functions

This section is a variation on [Sch95] in which we replace some properness hypotheses with that of being constructible up to infinity.

5.1 Correspondences

Consider the situation of Diagram (1.1) when replacing the manifolds X_i with b-analytic manifolds $X_{i\infty}$ ($i = 1, 2, 3$). Assume to be given two locally closed subsets subanalytic up to infinity:

$$S_1 \subset X_{12}, \quad S_2 \subset X_{23}.$$

We set, for $\varphi \in \mathcal{CF}(X_{1\infty})$

$$\mathcal{R}_{S_1}(\varphi) = \varphi \circ \mathbf{1}_{S_1} = \int_{q_2} q_1^* \varphi \cdot \mathbf{1}_{S_1}.$$

Set

$$(5.1) \quad \lambda := \mathbf{1}_{S_1} \circ \mathbf{1}_{S_2} \in \mathcal{CF}(X_{13\infty}).$$

Applying Theorem 4.14, we get that λ is well defined and moreover

$$(5.2) \quad \mathcal{R}_{S_2} \circ \mathcal{R}_{S_1}(\varphi) = \varphi \circ \lambda.$$

Now we assume that $X_1 = X_3$ and we change our notations, setting

$$X_1 = X_3 = X, \quad X_2 = Y.$$

For $(x, x') \in X \times X$, let

$$(5.3) \quad S_{12}(x, x') = \{y \in Y; (x, y) \in S_1, (y, x') \in S_2\} = (S_1 \times_Y S_2) \cap q_{13}^{-1}(x, x').$$

Then

$$(5.4) \quad \lambda(x, x') = \int_{q_{13}} \mathbf{1}_{S_1 \times_Y S_2} \cdot \mathbf{1}_{\{q_{13}^{-1}(x, x')\}} = \int_Y \mathbf{1}_{S_{12}(x, x')}.$$

We now consider the hypothesis

$$(5.5) \quad \begin{cases} \text{there exists } a, b \in \mathbb{Z} \text{ such that, for } (x, x') \in X \times X: \\ \lambda(x, x') = \begin{cases} a & \text{if } x \neq x', \\ b & \text{if } x = x'. \end{cases} \end{cases}$$

Writing $\lambda(x, x') = (b - a)\mathbf{1}_\Delta + a\mathbf{1}_{X \times X}$, we get:

Corollary 5.1 ([Sch95, Th. 3.1]). *Assume (5.5). Let $\varphi \in \mathcal{CF}(X)$. Then:*

$$\mathcal{R}_{S_2} \circ \mathcal{R}_{S_1}(\varphi) = (b - a)\varphi + a \int_X \varphi.$$

Here, $a \int_X \varphi \in \mathbb{Z}$ is identified with the constant function $(a \int_X \varphi) \cdot \mathbf{1}_X$.

Application to flag manifolds

Let \mathbb{W} be a real $(n+1)$ -dimensional vector space (with $n > 0$) and denote by $F_{n+1}(p, q)$, with $1 \leq p \leq q \leq n$, the set of pairs $\{(l, h)\}$ of linear subspaces of \mathbb{W} with $l \subset h$ and $\dim l = p$, $\dim h = q$. One sets $F_{n+1}(p) = F_{n+1}(p, p)$ and denotes as usual by q_1 and q_2 the two projections defined on $F_{n+1}(p) \times F_{n+1}(q)$. Then $F_{n+1}(p, q)$ is a real compact submanifold of $F_{n+1}(p) \times F_{n+1}(q)$, called the incidence relation. We denote by $F_{n+1}(q, p)$ its image by the map $F_{n+1}(p) \times F_{n+1}(q) \rightarrow F_{n+1}(q) \times F_{n+1}(p)$, $(x, y) \mapsto (y, x)$. In the sequel, we set

$$X = F_{n+1}(p), \quad Y = F_{n+1}(q), \quad S = F_{n+1}(p, q) \subset X \times Y, \quad S' = F_{n+1}(q, p) \subset Y \times X.$$

Now we shall assume $p = 1$ and $q > 1$. Recall that $F_{n+1}(1) = \mathbb{P}_n$, the n -dimensional real projective space.

In order to apply Corollary 5.1, it is enough to calculate $\lambda_{12}(x, x')$ given by (5.4) and (5.3) with $S_1 = S$ and $S_2 = S'$. Set

$$\mu_{n+1}(q) = \chi(F_{n+1}(q)).$$

Proposition 5.2. *Let $\varphi \in \mathcal{CF}(\mathbb{P}_n)$. Then:*

$$\mathcal{R}_{(n+1;q,1)} \circ \mathcal{R}_{(n+1;1,q)}(\varphi) = (\mu_n(q-1) - \mu_{n-1}(q-2))\varphi + \mu_{n-1}(q-2) \int_{\mathbb{P}_n} \varphi.$$

Proof. Let us represent x and x' by lines in \mathbb{W} and $y \in F_{n+1}(q)$ by a q -dimensional linear subspace. Then the set $S_{12}(x, x')$ is the set of q -dimensional linear subspaces of \mathbb{W} containing both the line x and the line x' . This set is isomorphic to $F_{n-1}(q-2)$ if $x \neq x'$ and to $F_n(q-1)$ if $x = x'$. \square

Of course, this formula is interesting only when $\mu_n(q-1) \neq \mu_{n-1}(q-2)$.

5.2 Application: the Radon transform

This section is extracted from [Sch95].

One can roughly describe the Radon transform as follows. How to reconstruct a function (say with compact support) on a real vector space \mathbb{V} from the knowledge of its integral along all affine hyperplanes? Since the family of these hyperplanes (including the hyperplane at infinity) is given by the dual projective space \mathbb{P}^* , where \mathbb{P} is the projective compactification of \mathbb{V} , it is natural to replace \mathbb{V} with \mathbb{P} .

We have $F_{n+1}(1) = \mathbb{P}_n$, the n -dimensional projective space and $F_{n+1}(n) = \mathbb{P}_n^*$, the dual projective space. The Radon transform thus corresponds to the case $p = 1$, $q = n$.

With the preceding notations, the incidence relation S is given by

$$S = F_{n+1}(1, n) = \{(x, y) \in \mathbb{P}_n \times \mathbb{P}_n^*; \langle x, y \rangle = 0\}.$$

The Radon transform of $\varphi \in \mathcal{CF}(\mathbb{P}_n)$, an element of $\mathcal{CF}(\mathbb{P}_n^*)$, is defined by

$$(5.6) \quad \mathcal{R}_{(n+1;1,n)}(\varphi) = \int_{\mathbb{P}_n} \mathbf{1}_S \cdot q_1^* \varphi = \varphi \circ \mathbf{1}_S.$$

For $y \in \mathbb{P}_n^*$, we shall denote by h_y its image in \mathbb{P}_n by the incidence relation:

$$h_y = \{x \in \mathbb{P}_n, \langle x, y \rangle = 0\}.$$

Therefore,

$$\mathcal{R}_{(n+1;1,n)}(\varphi)(y) = \int_{\mathbb{P}_n} \varphi \cdot \mathbf{1}_{h_y}.$$

Recall that the Euler-Poincaré index of \mathbb{P}_n is given by the formula:

$$(5.7) \quad \chi(\mathbb{P}_n) = \begin{cases} 1 & \text{if } n \text{ is even,} \\ 0 & \text{if } n \text{ is odd.} \end{cases}$$

Applying Proposition 5.2 together with (5.7), we get:

Corollary 5.3. *Let $\varphi \in \mathcal{CF}(\mathbb{P}_n)$. Then:*

$$\mathcal{R}_{(n+1,n,1)} \circ \mathcal{R}_{(n+1;1,n)}(\varphi) = \begin{cases} \varphi & \text{if } n \text{ is odd,} \\ -\varphi + \int_{\mathbb{P}_n} \varphi & \text{if } n \text{ is even and } n > 0. \end{cases}$$

Now assume $\dim \mathbb{V} = 3$ and let us calculate the Radon transform of the characteristic function $\mathbf{1}_K$ of a compact subanalytic subset K of \mathbb{V} (see (3.6)). First, consider a compact subanalytic subset L of a two dimensional affine vector space W . By Poincaré’s duality, there is an isomorphism $H_L^1(W; \mathbb{Q}_W) \simeq H^1(L; \mathbb{Q}_L)$ and moreover there is a short exact sequence:

$$0 \rightarrow H^0(W; \mathbb{Q}_W) \rightarrow H^0(W \setminus L; \mathbb{Q}_W) \rightarrow H_L^1(W; \mathbb{Q}_W) \rightarrow 0,$$

from which one deduces that:

$$b_1(L) = b_0(W \setminus L) - 1,$$

where b_i is the i -th Betti number. Note that $b_0(W \setminus L)$ is the number of connected components of $W \setminus L$, hence $b_1(L)$ is the “number of holes” of the compact set L . We may summarize:

Corollary 5.4. *The value at $y \in \mathbb{P}_3^*$ of the Radon transform of $\mathbf{1}_K$ is the number of connected components of $K \cap h_y$ minus the number of its holes.*

The inversion formula of the Radon transform tells us how to reconstruct the set K from the knowledge of the number of connected components and holes of all its affine slices.

References

- [BM88] E Bierstone and P. D. Milman, *Semi-analytic and subanalytic sets*, Publ. Math. I.H.E.S **67** (1988), 5-42.
- [CGR12] Justin Curry, Robert Ghrist, and Michael Robinson, *Euler Calculus with Applications to Signals and Sensing*, Proc. Sympos. Appl. Math., AMS (2012), available at [arXiv:1202.0275](https://arxiv.org/abs/1202.0275).
- [DK16] Andrea D’Agnolo and Masaki Kashiwara, *Riemann-Hilbert correspondence for holonomic D -modules*, Publ. IHES **123** (2016), 69-197.
- [VdD98] L. Van den Dries, *Tame Topology and O -minimal Structures*, Lect. Notes Series, vol. 248, London Math. Soc., 1998.
- [VdDM96] Lou Van den Dries and Chris Miller, *Geometric category and O -minimal Structures*, Duke math. Journ. **84** (1996), 497–539.
- [EP20] Mario Edmundo and Luca Prelli, *The six Grothendieck operations on o -minimal sheaves*, Math. Z **294**,n **1-2** (2020), 109–160, available at [arxiv:1401.0846v3](https://arxiv.org/abs/1401.0846v3).
- [Ern94] Lars Ernström, *Topological Radon transforms and the local Euler obstruction*, Duke Math. Journal **76** (1994), 1–21.
- [Gin86] Victor Ginsburg, *Characteristic varieties and vanishing cycles*, Inventiones Math. **84** (1986), 327–402.

- [Kas73] Masaki Kashiwara, *Index theorem for a maximally overdetermined system of linear differential equations*, Proc. Japan Acad. **49** (1973), 803–804.
- [Kas85] ———, *Index theorem for constructible sheaves*, Systèmes différentiels et singularités, Soc. Math. France, 1985, pp. 193–209.
- [KS90] Masaki Kashiwara and Pierre Schapira, *Sheaves on manifolds*, Grundlehren der Mathematischen Wissenschaften, vol. 292, Springer-Verlag, Berlin, 1990.
- [KS01] ———, *Ind-Sheaves*, Astérisque, vol. 271, Soc. Math. France, 2001.
- [KS16] ———, *Irregular holonomic kernels and Laplace transform*, Selecta Mathematica **22** (2016), 55–101.
- [KS18] ———, *Persistent homology and microlocal sheaf theory*, Journal of Applied and Computational Topology **2** (2018), 83–113, available at [arXiv:1705.00955](https://arxiv.org/abs/1705.00955).
- [KS19] ———, *Piecewise linear sheaves*, International Mathematics Research Notices (2019), available at [arXiv:math/1805.00349](https://arxiv.org/abs/math/1805.00349).
- [KS20] ———, *A finiteness theorem for holonomic DQ-modules on Poisson manifolds*, Tunisian Journ. Math. (2020), to appear, available at [arXiv:2001.06401](https://arxiv.org/abs/2001.06401).
- [McP74] Robert McPherson, *Chern classes of singular varieties*, Ann. Math. **100** (1974), 423–432.
- [Mil20a] Ezra Miller, *Homological algebra of modules over posets* (2020), available at [arXiv:2008.00063](https://arxiv.org/abs/2008.00063).
- [Mil20b] ———, *Stratifications of real vector spaces from constructible sheaves with conical micro-support* (2020), available at [arXiv:2008.00091](https://arxiv.org/abs/2008.00091).
- [Sab85] Claude Sabbah, *Quelques remarques sur la géométrie des espaces conormaux*, Astérisque **192** (1985), 161–192.
- [Sch89] Pierre Schapira, *Cycles Lagrangiens, fonctions constructibles et applications*, Sem EDP, Publ. Ec. Polyt. (1989).
- [Sch91] ———, *Operations on constructible functions*, Journ. Pure Appl. Algebra **72** (1991), 83–93.
- [Sch95] ———, *Tomography of constructible functions*, Vol. 948, Springer, Berlin, Applied algebra, algebraic algorithms and error-correcting codes, Lecture Notes in Comput. Sci., 1995, pp. 427–435.
- [Sch20] ———, *Constructible functions sheaves and functions up to infinity* (2020), available at [arXiv:2012.09652](https://arxiv.org/abs/2012.09652), v1, v2.
- [Sch03] Jörg Schürmann, *Topology of singular spaces and constructible sheaves*, Monografie Matematyczne, vol. 63, Springer Basel AG, 2003.
- [Vir88] Oleg Viro, *Some integral calculus based on Euler characteristic*, Vol. 1346, Springer, Berlin, Lecture Notes in Math, 1988, pp. 127–138.

Pierre Schapira
 Sorbonne Université, CNRS IMJ-PRG
 4 place Jussieu, 75252 Paris Cedex 05 France
 e-mail: pierre.schapira@imj-prg.fr
<http://webusers.imj-prg.fr/~pierre.schapira/>